



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 7/00	A2	(11) International Publication Number: WO 00/62744 (43) International Publication Date: 26 October 2000 (26.10.00)															
(21) International Application Number: PCT/US00/10490 (22) International Filing Date: 19 April 2000 (19.04.00) (30) Priority Data: <table border="0"> <tr> <td>60/129,975</td> <td>19 April 1999 (19.04.99)</td> <td>US</td> </tr> <tr> <td>09/441,303</td> <td>16 November 1999 (16.11.99)</td> <td>US</td> </tr> <tr> <td>60/175,315</td> <td>10 January 2000 (10.01.00)</td> <td>US</td> </tr> <tr> <td>09/541,965</td> <td>4 April 2000 (04.04.00)</td> <td>US</td> </tr> <tr> <td>09/544,783</td> <td>7 April 2000 (07.04.00)</td> <td>US</td> </tr> </table> (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors: ROBINSON, Larry, Richard; 1114 Tumbleweed Drive, Loveland, OH 45140 (US). BISSETT, Donald, Lynn; 3925 Dust Commander Drive, Hamilton, OH 45011 (US). DECKNER, George, Endel; 10572 Tanager Hills Drive, Cincinnati, OH 45249 (US). HA, Robert, Bao, Kim; 1199 Mellie Avenue, Milford, OH 45150 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).		60/129,975	19 April 1999 (19.04.99)	US	09/441,303	16 November 1999 (16.11.99)	US	60/175,315	10 January 2000 (10.01.00)	US	09/541,965	4 April 2000 (04.04.00)	US	09/544,783	7 April 2000 (07.04.00)	US	(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
60/129,975	19 April 1999 (19.04.99)	US															
09/441,303	16 November 1999 (16.11.99)	US															
60/175,315	10 January 2000 (10.01.00)	US															
09/541,965	4 April 2000 (04.04.00)	US															
09/544,783	7 April 2000 (07.04.00)	US															
(54) Title: SKIN CARE COMPOSITIONS CONTAINING COMBINATION OF SKIN CARE ACTIVES																	
(57) Abstract The present invention relates to skin care compositions containing combinations of skin care actives and to methods of using such compositions to regulate the condition of skin. The compositions contain a safe and effective amount of farnesol; a safe and effective amount of bisabolol; a safe and effective amount of at least one additional skin care active; and a dermatologically acceptable carrier.																	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SKIN CARE COMPOSITIONS CONTAINING COMBINATION
OF SKIN CARE ACTIVES

TECHNICAL FIELD

The present invention relates to topical compositions containing skin care actives, particularly farnesol, bisabolol and at least one additional skin care active. Such compositions are useful for regulating the condition of skin, especially for regulating visible and/or tactile discontinuities in skin associated, e.g., with skin aging. Preferred compositions also contain phytantriol.

CROSS REFERENCE

This application is a continuation-in-part of a co-pending application, (no application number received yet) P&G Case No. 7525M, filed on April 4, 2000, which in turn claims priority under Title 35, United States Code § 119(e) from Provisional Application Serial No. 60/129,975, filed on April 19, 1999. This application is also a continuation-in-part of U.S. Application Serial No. 09/441,303, filed November 16, 1999. Furthermore, this application claims priority under Title 35, United States Code § 119(e) from Provisional Application Serial No. 60/175,315, filed January 10, 2000.

CROSS REFERENCE

This application is a continuation-in-part of U.S. Application Serial No. 09/441,303, filed November 16, 1999. This application is a continuation-in-part of a co-pending application, (no application number received yet) P&G Case No. 7525M, filed on April 4, 2000, which in turn claims priority under Title 35, United States Code § 119(e) from Provisional Application Serial No. 60/129,975, filed on April 19, 1999. This application also claims priority under Title 35, United States Code § 119(e) from Provisional Application Serial No. 60/175,315 filed January 10, 2000.

BACKGROUND

Many personal care products currently available to consumers are directed primarily to improving the health and/or physical appearance of the skin. Among these skin care products, many are directed to delaying, minimizing or even eliminating skin wrinkling and other histological changes typically associated with the aging of skin or environmental damage to human skin. Numerous compounds have been described in the art as being useful for regulating skin condition, including regulating fine lines, wrinkles and other forms of uneven or rough surface texture associated with aged or photodamaged skin.

Skin is subject to insults by many extrinsic and intrinsic factors. Extrinsic factors include ultraviolet radiation (e.g., from sun exposure), environmental pollution, wind, heat, low humidity, harsh surfactants, abrasives, and the like. Intrinsic factors include chronological aging and other biochemical changes from within the skin. Whether extrinsic or intrinsic, these factors result in visible signs of skin aging and environmental damage, such as wrinkling and other forms of roughness (including increased pore size, flaking and skin lines), and other histological changes associated with skin aging or damage. To many people, skin wrinkles are a reminder of the disappearance of youth. As a result, the elimination of wrinkles has become a booming business in youth-conscious societies. Treatments range from cosmetic creams and moisturizers to various forms of cosmetic surgery.

Extrinsic or intrinsic factors may result in the thinning and general degradation of the skin. For example, as the skin naturally ages, there is a reduction in the cells and blood vessels that supply the skin. There is also a flattening of the dermal-epidermal junction which results in weaker mechanical resistance of this junction. See, for example, Oikarinen, "The Aging of Skin: Chronoaging Versus Photoaging," *Photodermatol. Photoimmunol. Photomed.*, vol. 7, pp. 3-4, 1990, which is incorporated by reference herein in its entirety.

A large number of skin care actives are known in the art and used to improve the health and/or physical appearance of the skin. For example, salicylic acid and benzoyl peroxide are used in skin care compositions to treat acne. Retinoids are another example of skin care actives used in skin care compositions to reduce signs of aging skin. Although formulating skin care compositions with such actives provide skin care benefits, there are also challenges in formulating such compositions. For example, retinoid compositions typically have to be prepared under specialized conditions, such as in an inert atmosphere, and may exhibit less than optimal stability, such as discoloration, at times. Some skin care active containing compositions may result in skin irritation, such as stinging, burning, and redness.

Based on the foregoing, there is a continuing need to formulate skin care compositions which improve the health and/or physical appearance of the skin, which are for example, aesthetically pleasing, stable, and effective in treating the appearance of wrinkles, fine lines, pores, poor skin color (e.g. redness, sallowness, and other forms of undesirable skin surface texture).

Surprisingly, it has now been found that compositions containing a multiple active system, particularly farnesol in combination with bisabolol and at least one additional skin care active, provide benefits in regulating skin condition previously unrecognized in the art of which

the present inventors are aware. For example, topical applications of farnesol in combination with bisabolol and at least one additional skin care active synergistically regulate (prophylactically and/or therapeutically) visible and/or tactile discontinuities in mammalian skin, including fine lines, wrinkles, enlarged pores, roughness, dryness, and other skin texture discontinuities, e.g., reduces or effaces the visibility of fine lines, wrinkles, and other forms of uneven or rough surface texture associated with aged or photodamaged skin.

None of the existing art provides all of the advantages and benefits of the present invention.

SUMMARY

The present invention relates to a composition containing a safe and effective amount of farnesol; a safe and effective amount of bisabolol; a safe and effective amount of at least one additional skin care active; and a dermatologically acceptable carrier.

The present invention also relates to methods of using such compositions to regulate the condition of mammalian skin. Said methods generally contain the step of topically applying the composition to the skin of a mammal needing such treatment, a safe and effective amount of such compositions.

Preferred embodiments include phytantriol.

These and other features, aspects, and advantages of the present invention will become evident to those skilled in the art from a reading of the present disclosure.

DETAILED DESCRIPTION

While the specification concludes with the claims particularly pointing and distinctly claiming the invention, it is believed that the present invention will be better understood from the following description.

All percentages and ratios used herein are by weight of the total composition and all measurements made are at 25°C, unless otherwise designated.

The compositions of the present invention can comprise, consist essentially of, or consist of, the components of the present invention as well as other ingredients described herein. As used herein, "consisting essentially of" means that the composition or component may include additional ingredients, but only if the additional ingredients do not materially alter the basic and novel characteristics of the claimed compositions or methods.

All publications cited herein are hereby incorporated by reference in their entirety.

The term "keratinous tissue," as used herein, refers to keratin-containing layers disposed as the outermost protective covering of mammals (e.g., humans, dogs, cats, etc.) which includes, but is not limited to, skin, lips, hair, toenails, fingernails, cuticles, hooves, etc.

The term "topical application", as used herein, means to apply or spread the compositions of the present invention onto the surface of the keratinous tissue.

The term "dermatologically-acceptable," as used herein, means that the compositions or components thereof so described are suitable for use in contact with mammalian keratinous tissue without undue toxicity, incompatibility, instability, allergic response, and the like.

The term "safe and effective amount" as used herein means an amount of a compound or composition sufficient to significantly induce a positive benefit, preferably a positive keratinous tissue appearance or feel benefit, including independently or in combinations the benefits disclosed herein, but low enough to avoid serious side effects, i.e., to provide a reasonable benefit to risk ratio, within the scope of sound judgment of the skilled artisan.

The term "sagging" as used herein means the laxity, slackness, or the like condition of skin that occurs as a result of loss of, damage to, alterations to, and/or abnormalities in dermal elastin.

The terms "smoothing" and "softening" as used herein mean altering the surface of the keratinous tissue such that its tactile feel is improved.

"Signs of skin aging" include, but are not limited to, all outward visibly and tactilely perceptible manifestations as well as any other macro or micro effects due to skin aging. Such signs may be induced or caused by intrinsic factors or extrinsic factors, e.g., chronological aging and/or environmental damage. These signs may result from processes which include, but are not limited to, the development of textural discontinuities such as wrinkles and coarse deep wrinkles, skin lines, crevices, bumps, large pores (e.g., associated with adnexal structures such as sweat gland ducts, sebaceous glands, or hair follicles), or unevenness or roughness, loss of skin elasticity (loss and/or inactivation of functional skin elastin), sagging (including puffiness in the eye area and jowls), loss of skin firmness, loss of skin tightness, loss of skin recoil from deformation, discoloration (including undereye circles), blotching, sallowness, hyperpigmented skin regions such as age spots and freckles, keratoses, abnormal differentiation, hyperkeratinization, elastosis, collagen breakdown, and other histological changes in the stratum corneum, dermis, epidermis, the skin vascular system (e.g., telangiectasia or spider vessels), and underlying tissues, especially those proximate to the skin.

The present invention is useful for therapeutically regulating visible and/or tactile discontinuities in mammalian skin, including discontinuities in skin texture and color. For example, the apparent diameter of pores decreases, the apparent height of tissue immediately proximate to pore openings approaches that of the interadnexal skin, the skin tone/color becomes more uniform, and/or the length, depth, and/or other dimension of lines and/or wrinkles are decreased.

The compositions of the present invention are also useful for regulating the condition of skin and especially for regulating keratinous tissue condition. Regulation of skin condition, namely mammalian and in particular human skin condition, is often required due to conditions which may be induced or caused by factors internal and/or external to the body. Examples include, environmental damage, radiation exposure (including ultraviolet radiation), chronological aging, menopausal status (e.g., post-menopausal changes in skin), stress, diseases, etc. For instance, "regulating skin condition" includes prophylactically regulating and/or therapeutically regulating skin condition, and may involve one or more of the following benefits: thickening of skin (i.e., building the epidermis and/or dermis and/or sub-dermal (e.g., subcutaneous fat or muscle) layers of the skin and where applicable the keratinous layers of the nail and hair shaft) to reduce skin atrophy, increasing the convolution of the dermal-epidermal border (also known as the rete ridges), preventing loss of skin elasticity (loss, damage and/or inactivation of functional skin elastin) such as elastosis, sagging, loss of skin recoil from deformation; non-melanin skin discoloration such as under eye circles, blotching (e.g., uneven red coloration due to, e.g., rosacea) (hereinafter referred to as "red blotchiness"), sallowness (pale color), discoloration caused by telangiectasia or spider vessels.

As used herein, prophylactically regulating skin condition includes delaying, minimizing and/or preventing visible and/or tactile discontinuities in skin (e.g., texture irregularities in the skin which may be detected visually or by feel).

As used herein, therapeutically regulating skin condition includes ameliorating, e.g., diminishing, minimizing and/or effacing, discontinuities in skin.

The compositions of the present invention are also useful for improving skin appearance and/or feel. For example, compositions of the present invention are useful for regulating the appearance of skin condition by providing an immediate visual improvement in skin appearance following application of the composition to the skin. Generally speaking, compositions of the present invention which further contain particulate materials will be most useful for providing the immediate visual improvement.

The compositions of the present invention provide additional benefits, including stability, absence of significant (consumer-unacceptable) skin irritation and good aesthetics.

The compositions of the present invention are stable. The ingredients used herein, including the farnesol, bisabolol, and other skin care actives are stable in the composition and are compatible with each other. Therefore, the compositions containing the combination of farnesol, bisabolol, and at least one additional skin care active are capable of providing additive and/or synergistic skin benefits. Additionally, the resulting skin care composition has good product stability and a reasonably long shelf-life.

The resulting compositions containing combinations of farnesol, bisabolol, and at least one additional skin care active have good aesthetics. Examples of good aesthetics include compositions, such as luxurious creams and lotions, that (i) are light and nongreasy, (ii) have a smooth, silky feel upon the skin, (iii) spread easily, and/or (iv) absorb quickly. Other examples of good aesthetics include compositions that have a consumer acceptable appearance (i.e. no unpleasant odor or discoloration present), and provide good skin feel.

The compositions of the present invention contain farnesol, bisabolol, at least one additional skin care active, and a dermatologically acceptable carrier. The compositions herein may also include a wide variety of optional ingredients.

The compositions of the present invention, are described in detail hereinafter.

I. Farnesol

The topical compositions of the present invention contain a safe and effective amount of farnesol. Farnesol is a naturally occurring substance which is believed to act as a precursor and/or intermediate in the biosynthesis of squalene and sterols, especially cholesterol. Farnesol is also involved in protein modification and regulation (e.g., farnesylation of proteins), and there is a cell nuclear receptor which is responsive to farnesol.


Chemically, farnesol is [2E,6E]-3,7,11-trimethyl-2,6,10-dodecatrien-1-ol and as used herein "farnesol" includes isomers and tautomers of such. Farnesol is commercially available, e.g., under the names farnesol (a mixture of isomers from Dragoco, 10 Gordon Drive, Totowa, New Jersey) and trans-trans-farnesol (Sigma Chemical Company, P. O. Box 14508, St. Louis, Missouri).

The compositions herein preferably contain from about 0.001% to about 50%, by weight of the composition, more preferably from about 0.01% to about 20%, even more preferably from about 0.1% to about 15%, even more preferably from about 0.1% to about 10%, still more

preferably from about 0.5% to about 5%, and still more preferably from about 1% to about 5% of farnesol.

II. Bisabolol

The topical compositions of the present invention contain a safe and effective amount of bisabolol. Bisabolol is a naturally occurring unsaturated monocyclic terpene alcohol having the following structure



It is the primary active component of chamomile extract/oil. Bisabolol can be synthetic (d,l - alpha-isomer or (+/-)-alpha-isomer) or natural ((-)-alpha-isomer) in origin and can be used as essentially pure compounds or mixtures of compounds (e.g., extracts from natural sources such as chamomile). The alpha form of bisabolol (α -bisabolol) is used in a variety of cosmetic products as a skin conditioning or soothing agent. As used herein, "bisabolol" includes chamomile extract or oil and any isomers and tautomers of such. Suitable bisabolol compounds are commercially available as a natural material from Dragoco (Totowa, New Jersey) under the product name alpha-bisabolol natural and as a synthetic material from Fluka (Milwaukee, Wisconsin) under the product name alpha-bisabolol.

In the compositions of the present invention, the composition preferably contains from about 0.001% to about 50%, by weight of the composition, more preferably from about 0.01% to about 20%, even more preferably from about 0.01% to about 15%, and still more preferably from about 0.1% to about 10%, even more preferably from about 0.1% to about 5%.

III. Additional Skin Care Active

The compositions of the present invention contain at least one additional skin care active. The compositions of the present invention may contain additional skin care actives as well.

In a preferred embodiment, where the composition is to be in contact with human keratinous tissue, the additional skin care active(s) should be suitable for application to keratinous tissue, that is, when incorporated into the composition they are suitable for use in contact with human keratinous tissue without undue toxicity, incompatibility, instability, allergic

response, and the like within the scope of sound medical judgment. The *CTFA Cosmetic Ingredient Handbook*, Second Edition (1992) describes a wide variety of nonlimiting cosmetic and pharmaceutical ingredients commonly used in the skin care industry, which are suitable for use in the compositions of the present invention. Examples of these ingredient classes include: abrasives, absorbents, aesthetic components such as fragrances, pigments, colorings/colorants, essential oils, skin sensates, astringents, etc. (e.g., clove oil, menthol, camphor, eucalyptus oil, eugenol, menthyl lactate, witch hazel distillate), anti-acne agents, anti-caking agents, antifoaming agents, antimicrobial agents (e.g., iodopropyl butylcarbamate), antioxidants, binders, biological additives, buffering agents, bulking agents, chelating agents, chemical additives, colorants, cosmetic astringents, cosmetic biocides, denaturants, drug astringents, external analgesics, film formers or materials, e.g., polymers, for aiding the film-forming properties and substantivity of the composition (e.g., copolymer of eicosene and vinyl pyrrolidone), opacifying agents, pH adjusters, propellants, reducing agents, sequestrants, skin bleaching and lightening agents (e.g., hydroquinone, kojic acid, ascorbic acid, magnesium ascorbyl phosphate, ascorbyl glucosamine), skin-conditioning agents (e.g., humectants, including miscellaneous and occlusive), skin soothing and/or healing agents (e.g., panthenol and derivatives (e.g., ethyl panthenol), aloe vera, pantothenic acid and its derivatives, allantoin, bisabolol, and dipotassium glycyrrhizinate), skin treating agents, thickeners, and vitamins and derivatives thereof.

In any embodiment of the present invention, however, the actives useful herein can be categorized by the benefit they provide or by their postulated mode of action. However, it is to be understood that the actives useful herein can in some instances provide more than one benefit or operate via more than one mode of action. Therefore, classifications herein are made for the sake of convenience and are not intended to limit the active to that particular application or applications listed.

Phytantriol

The topical compositions of the present invention may contain a safe and effective amount of phytantriol. Phytantriol is the common name for the chemical known as 3,7,11,15-tetramethylhexadecane-1,2,3,-triol. Phytantriol is commercially available from BASF (1609 Biddle Avenue, Whyandotte, MI). For example, phytantriol is useful as a spider vessel/red blotchiness repair agent, a dark circle/puffy eye repair agent, sallowness repair agent, a sagging repair agent, an anti-itch agent, a skin thickening agent, a pore reduction agent, oil/shine

reduction agent, a post-inflammatory hyperpigmentation repair agent, wound treating agent, an anti-cellulite agent, and regulating skin texture, including wrinkles and fine lines.

When included in the compositions of the present invention, the phytantriol preferably is included in an amount from about 0.001% to about 50% by weight of the composition, more preferably from about 0.01% to about 20%, even more preferably from about 0.1% to about 15%, even more preferably from about 0.2% to about 10%, still more preferably from about 0.5% to about 10%, and still more preferably from about 1% to about 5%.

Desquamation Actives

A safe and effective amount of a desquamation active may be added to the compositions of the present invention. more preferably from about 0.1% to about 10%, even more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 4%, by weight of the composition. Desquamation actives enhance the skin appearance benefits of the present invention. For example, the desquamation actives tend to improve the texture of the skin (e.g., smoothness). One desquamation system that is suitable for use herein contains sulfhydryl compounds and zwitterionic surfactants and is described in U.S. Patent No. 5,681,852, to Bissett, incorporated herein by reference. Another desquamation system that is suitable for use herein contains salicylic acid and zwitterionic surfactants and is described in U.S. Patent No. 5,652,228 to Bissett, incorporated herein by reference. Zwitterionic surfactants such as described in these applications are also useful as desquamatory agents herein, with cetyl betaine being particularly preferred.

Anti-Acne Actives

The compositions of the present invention may contain a safe and effective amount of one or more anti-acne actives. Examples of useful anti-acne actives include resorcinol, sulfur, salicylic acid, benzoyl peroxide, erythromycin, zinc, etc. Further examples of suitable anti-acne actives are described in further detail in U. S. Patent No. 5,607,980, issued to McAtee et al, on March 4, 1997.

Anti-Wrinkle Actives/Anti-Atrophy Actives


The compositions of the present invention may further contain a safe and effective amount of one or more anti-wrinkle actives or anti-atrophy actives. Exemplary anti-wrinkle/anti-atrophy actives suitable for use in the compositions of the present invention include sulfur-containing D and L amino acids and their derivatives and salts, particularly the N-acetyl derivatives, a preferred example of which is N-acetyl-L-cysteine; thiols, e.g. ethane thiol; hydroxy acids (e.g., alpha-hydroxy acids such as lactic acid and glycolic acid or beta-hydroxy

acids such as salicylic acid and salicylic acid derivatives such as the octanoyl derivative), phytic acid, lipoic acid; lysophosphatidic acid, skin peel agents (e.g., phenol and the like), vitamin B₃ compounds and retinoids which enhance the keratinous tissue appearance benefits of the present invention, especially in regulating keratinous tissue condition, e.g., skin condition.

a) Vitamin B₃ Compounds

The compositions of the present invention may contain a safe and effective amount of a vitamin B₃ compound. Vitamin B₃ compounds are particularly useful for regulating skin condition as described in co-pending U. S. Application Serial No. 08/834,010, filed April 11, 1997 (corresponding to international publication WO 97/39733 A1, published October 30, 1997). When vitamin B₃ compounds are present in the compositions of the instant invention, the compositions preferably contain from about 0.01% to about 50%, more preferably from about 0.1% to about 10%, even more preferably from about 0.5% to about 10%, and still more preferably from about 1% to about 5%, still more preferably from about 2% to about 5%, by weight of the composition, of the vitamin B₃ compound.

As used herein, "vitamin B₃ compound" means a compound having the formula:



wherein R is - CONH₂ (i.e., niacinamide), - COOH (i.e., nicotinic acid) or - CH₂OH (i.e., nicotinyl alcohol); derivatives thereof; and salts of any of the foregoing.

Exemplary derivatives of the foregoing vitamin B₃ compounds include nicotinic acid esters, including non-vasodilating esters of nicotinic acid (e.g., tocopheryl nicotinate), nicotinyl amino acids, nicotinyl alcohol esters of carboxylic acids, nicotinic acid N-oxide and niacinamide N-oxide.

Examples of suitable vitamin B₃ compounds are well known in the art and are commercially available from a number of sources, e.g., the Sigma Chemical Company (St. Louis, MO); ICN Biomedicals, Inc. (Irvin, CA) and Aldrich Chemical Company (Milwaukee, WI).

The vitamin compounds may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources.

b) Retinoids

The compositions of the present invention may also contain a retinoid. As used herein, "retinoid" includes all natural and/or synthetic analogs of Vitamin A or retinol-like compounds which possess the biological activity of Vitamin A in the skin as well as the geometric isomers and stereoisomers of these compounds. The retinoid is preferably retinol, retinol esters (e.g., C₂ - C₂₂ alkyl esters of retinol, including retinyl palmitate, retinyl acetate, retinyl propionate), retinal, and/or retinoic acid (including all-trans retinoic acid and/or 13-cis-retinoic acid), more preferably retinoids other than retinoic acid. These compounds are well known in the art and are commercially available from a number of sources, e.g., Sigma Chemical Company (St. Louis, MO), and Boehringer Mannheim (Indianapolis, IN). Other retinoids which are useful herein are described in U.S. Patent Nos. 4,677,120, issued Jun. 30, 1987 to Parish et al.; 4,885,311, issued Dec. 5, 1989 to Parish et al.; 5,049,584, issued Sep. 17, 1991 to Purcell et al.; 5,124,356, issued Jun. 23, 1992 to Purcell et al.; and Reissue 34,075, issued Sep. 22, 1992 to Purcell et al.. Other suitable retinoids are tocopheryl-retinoate [tocopherol ester of retinoic acid (trans- or cis-), adapalene {6-[3-(1-adamantyl)-4-methoxyphenyl]-2-naphthoic acid}, and tazarotene (ethyl 6-[2-(4,4-dimethylthiochroman-6-yl)-ethynyl]nicotinate). Preferred retinoids are retinol, retinyl palmitate, retinyl acetate, retinyl propionate, retinal and combinations thereof.

The retinoid may be included as the substantially pure material, or as an extract obtained by suitable physical and/or chemical isolation from natural (e.g., plant) sources. The retinoid is preferably substantially pure, more preferably essentially pure.

The compositions of this invention may contain a safe and effective amount of the retinoid, such that the resultant composition is safe and effective for regulating keratinous tissue condition, preferably for regulating visible and/or tactile discontinuities in skin, more preferably for regulating signs of skin aging, even more preferably for regulating visible and/or tactile discontinuities in skin texture associated with skin aging. The compositions preferably contain from or about 0.005% to or about 2%, more preferably 0.01% to or about 2%, retinoid. Retinol is preferably used in an amount of from or about 0.01% to or about 0.15%; retinol esters are preferably used in an amount of from or about 0.01% to or about 2% (e.g., about 1%); retinoic acids are preferably used in an amount of from or about 0.01% to or about 0.25%; tocopheryl-retinoate, adapalene, and tazarotene are preferably used in an amount of from or about 0.01% to or about 2%.

Where the compositions of the present invention contain both a retinoid and a Vitamin B₃ compound, the retinoid is preferably used in the above amounts, and the vitamin B₃

compound is preferably used in an amount of from or about 0.1% to or about 10%, more preferably from or about 2% to or about 5%.

(c) Hydroxy Acids

The compositions of the present invention may contain a safe and effective amount of a Hydroxy Acid. Preferred hydroxy acids for use in the compositions of the present invention include salicylic acid and salicylic acid derivatives. When present in the compositions of the present invention, salicylic acid is preferably used in an amount of from about 0.01% to about 50%, more preferably from about 0.1% to about 20%, even more preferably from about 0.1% to about 10%, still more preferably from about 0.5% to about 5%, and still more preferably from about 0.5% to about 2%.

Peptides

Peptides, including but not limited to, di-, tri-, tetra-, and pentapeptides and derivatives thereof, may be included in the compositions of the present invention in amounts that are safe and effective. As used herein, "peptides" refers to both the naturally occurring peptides and synthesized peptides. Also useful herein are naturally occurring and commercially available compositions that contain peptides.

Suitable dipeptides for use herein include Carnosine (beta-ala-his). Suitable tripeptides for use herein include, gly-his-lys, arg-lys-arg, his-gly-gly. Preferred tripeptides and derivatives thereof include palmitoyl-gly-his-lys, which may be purchased as Biopeptide CL® (100ppm of palmitoyl-gly-his-lys commercially available from Sederma, France); Peptide CK (arg-lys-arg); Peptide CK+ (ac-arg-lys-arg-NH₂); and a copper derivative of his-gly-gly sold commercially as Iamin, from Sigma (St.Louis, Missouri). Suitable tetrapeptides for use herein include Peptide E, arg-ser-arg-lys (SEQ ID NO:1). Suitable pentapeptides for use herein include lys-thr-thr-lys-ser. A preferred commercially available pentapeptide derivative composition is Matrixyl®, which contains 100 ppm palmitoyl-lys-thr-thr-lys-ser (SEQ ID NO:2, commercially available from Sederma, France).

Preferably, the peptide is selected from palmitoyl-lys-thr-thr-lys-ser, palmitoyl-gly-his-lys, beta-ala-his, their derivatives, and combinations thereof. More preferably, the peptide is selected from palmitoyl-lys-thr-thr-lys-ser, palmitoyl-gly-his-lys, their derivatives, and combinations thereof. Even more preferably, the peptide is selected from palmitoyl-lys-thr-thr-lys-ser and derivatives thereof.

When included in the present compositions, peptides are preferably included in amounts of from about $1 \times 10^{-6}\%$ to about 10%, more preferably from about $1 \times 10^{-6}\%$ to about 0.1%, even

more preferably from about $1 \times 10^{-5}\%$ to about 0.01%, by weight of the composition. In certain compositions where the peptide is Carnosine®, the compositions preferably contain from about 0.1% to about 5%, by weight of the composition, of such peptides. In other embodiments wherein the peptide-containing compositions, Matrixyl®, and/or Biopeptide CL® are included, the compositions preferably contain from about 0.1% to about 10%, by weight compositions, of Matrixyl® and/or Biopeptide CL® peptide-containing compositions.

Anti-Oxidants/Radical Scavengers

The compositions of the present invention may include a safe and effective amount of an anti-oxidant/radical scavenger. The anti-oxidant/radical scavenger is especially useful for providing protection against UV radiation which can cause increased scaling or texture changes in the stratum corneum and against other environmental agents which can cause skin damage.

A safe and effective amount of an anti-oxidant/radical scavenger may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition.

Anti-oxidants/radical scavengers such as ascorbic acid (vitamin C) and its salts, ascorbyl esters of fatty acids, ascorbic acid derivatives (e.g., magnesium ascorbyl phosphate, sodium ascorbyl phosphate, ascorbyl sorbate), tocopherol (vitamin E), tocopherol sorbate, tocopherol acetate, other esters of tocopherol, butylated hydroxy benzoic acids and their salts, 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (commercially available under the tradename Trolox®), gallic acid and its alkyl esters, especially propyl gallate, uric acid and its salts and alkyl esters, sorbic acid and its salts, lipoic acid, amines (e.g., N,N-diethylhydroxylamine, amino-guanidine), sulfhydryl compounds (e.g., glutathione), dihydroxy fumaric acid and its salts, lysine pidolate, arginine pilolate, nordihydroguaiaretic acid, bioflavonoids, curcumin, lysine, methionine, proline, superoxide dismutase, silymarin, tea extracts, grape skin/seed extracts, melanin, and rosemary extracts may be used. Preferred anti-oxidants/radical scavengers are selected from tocopherol sorbate and other esters of tocopherol, more preferably tocopherol sorbate. For example, the use of tocopherol sorbate in topical compositions and applicable to the present invention is described in U.S. Patent No. 4,847,071, issued on July 11, 1989 to Donald L. Bissett, Rodney D. Bush and Ranjit Chatterjee.

Chelators

The compositions of the present invention may also contain a safe and effective amount of a chelator or chelating agent. As used herein, "chelator" or "chelating agent" means an active agent capable of removing a metal ion from a system by forming a complex so that the metal ion

cannot readily participate in or catalyze chemical reactions. The inclusion of a chelating agent is especially useful for providing protection against UV radiation which can contribute to excessive scaling or skin texture changes and against other environmental agents which can cause skin damage.

A safe and effective amount of a chelating agent may be added to the compositions of the subject invention, preferably from about 0.1% to about 10%, more preferably from about 1% to about 5%, of the composition. Exemplary chelators that are useful herein are disclosed in U.S. Patent No. 5,487,884, issued 1/30/96 to Bissett et al.; International Publication No. 91/16035, Bush et al., published 10/31/95; and International Publication No. 91/16034, Bush et al., published 10/31/95. Preferred chelators useful in compositions of the subject invention are furildioxime, furilmonoxime, and derivatives thereof.

Flavonoids

The compositions of the present invention may optionally contain a flavonoid compound. Flavonoids are broadly disclosed in U.S. Patents 5,686,082 and 5,686,367, both of which are herein incorporated by reference. Flavonoids suitable for use in the present invention are flavanones selected from unsubstituted flavanones, mono-substituted flavanones, and mixtures thereof; chalcones selected from unsubstituted chalcones, mono-substituted chalcones, di-substituted chalcones, tri-substituted chalcones, and mixtures thereof; flavones selected from unsubstituted flavones, mono-substituted flavones, di-substituted flavones, and mixtures thereof; one or more isoflavones; coumarins selected from unsubstituted coumarins, mono-substituted coumarins, di-substituted coumarins, and mixtures thereof; chromones selected from unsubstituted chromones, mono-substituted chromones, di-substituted chromones, and mixtures thereof; one or more dicoumarols; one or more chromanones; one or more chromanols; isomers (e.g., cis/trans isomers) thereof; and mixtures thereof. By the term "substituted" as used herein means flavonoids wherein one or more hydrogen atom of the flavonoid has been independently replaced with hydroxyl, C1-C8 alkyl, C1-C4 alkoxy, O-glycoside, and the like or a mixture of these substituents.

Examples of suitable flavonoids include, but are not limited to, unsubstituted flavanone, mono-hydroxy flavanones (e.g., 2'-hydroxy flavanone, 6-hydroxy flavanone, 7-hydroxy flavanone, etc.), mono-alkoxy flavanones (e.g., 5-methoxy flavanone, 6-methoxy flavanone, 7-methoxy flavanone, 4'-methoxy flavanone, etc.), unsubstituted chalcone (especially unsubstituted trans-chalcone), mono-hydroxy chalcones (e.g., 2'-hydroxy chalcone, 4'-hydroxy chalcone, etc.), di-hydroxy chalcones (e.g., 2',4'-dihydroxy chalcone, 2',4'-dihydroxy chalcone,

2,2'-dihydroxy chalcone, 2',3-dihydroxy chalcone, 2',5'-dihydroxy chalcone, etc.), and trihydroxy chalcones (e.g., 2',3',4'-trihydroxy chalcone, 4,2',4'-trihydroxy chalcone, 2,2',4'-trihydroxy chalcone, etc.), unsubstituted flavone, 7,2'-dihydroxy flavone, 3',4'-dihydroxy naphthoflavone, 4'-hydroxy flavone, 5,6-benzoflavone, and 7,8-benzoflavone, unsubstituted isoflavone, daidzein (7,4'-dihydroxy isoflavone), 5,7-dihydroxy-4'-methoxy isoflavone, soy isoflavones (a mixture extracted from soy), unsubstituted coumarin, 4-hydroxy coumarin, 7-hydroxy coumarin, 6-hydroxy-4-methyl coumarin, unsubstituted chromone, 3-formyl chromone, 3-formyl-6-isopropyl chromone, unsubstituted dicoumarol, unsubstituted chromanone, unsubstituted chromanol, and mixtures thereof..

Preferred for use herein are unsubstituted flavanone, methoxy flavanones, unsubstituted chalcone, 2',4-dihydroxy chalcone, and mixtures thereof. More preferred are unsubstituted flavanone, unsubstituted chalcone (especially the trans isomer), and mixtures thereof.

They can be synthetic materials or obtained as extracts from natural sources (e.g., plants). The naturally sourced material can also further be derivatized (e.g., an ester or ether derivative prepared following extraction from a natural source). Flavonoid compounds useful herein are commercially available from a number of sources, e.g., Indofine Chemical Company, Inc. (Somerville, New Jersey), Steraloids, Inc. (Wilton, New Hampshire), and Aldrich Chemical Company, Inc. (Milwaukee, Wisconsin).

Mixtures of the above flavonoid compounds may also be used.

The herein described flavonoid compounds are preferably present in the instant invention at concentrations of from about 0.01% to about 20%, more preferably from about 0.1% to about 10% , and still more preferably from about 0.5% to about 5%.

Anti-Inflammatory Agents

A safe and effective amount of an anti-inflammatory agent may be added to the compositions of the present invention, preferably from about 0.1% to about 10%, more preferably from about 0.5% to about 5%, of the composition. The anti-inflammatory agent enhances the skin appearance benefits of the present invention, e.g., such agents contribute to a more uniform and acceptable skin tone or color. The exact amount of anti-inflammatory agent to be used in the compositions will depend on the particular anti-inflammatory agent utilized since such agents vary widely in potency.

Steroidal anti-inflammatory agents, including but not limited to, corticosteroids such as hydrocortisone, hydroxyltriamcinolone, alpha-methyl dexamethasone, dexamethasone-phosphate, beclomethasone dipropionates, clobetasol valerate, desonide, desoxymethasone,

desoxycorticosterone acetate, dexamethasone, dichlorisone, diflorasone diacetate, diflucortolone valerate, fluadrenolone, fluclorolone acetonide, fludrocortisone, flumethasone pivalate, fluosinolone acetonide, fluocinonide, flucortine butylesters, fluocortolone, fluprednidene (fluprednylidene) acetate, flurandrenolone, halcinonide, hydrocortisone acetate, hydrocortisone butyrate, methylprednisolone, triamcinolone acetonide, cortisone, cortodoxone, flucetonide, fludrocortisone, difluorosone diacetate, fluradrenolone, fludrocortisone, difluorosone diacetate, fluradrenolone acetonide, medrysone, amcinafel, amcinafide, betamethasone and the balance of its esters, chloroprednisone, chlorprednisone acetate, clocortelone, clescinsolone, dichlorisone, diflurprednate, flucloronide, flunisolide, fluoromethalone, fluperolone, fluprednisolone, hydrocortisone valerate, hydrocortisone cyclopentylpropionate, hydrocortamate, meprednisone, paramethasone, prednisolone, prednisone, beclomethasone dipropionate, triamcinolone, and mixtures thereof may be used. The preferred steroidal anti-inflammatory for use is hydrocortisone.

A second class of anti-inflammatory agents which is useful in the compositions includes the nonsteroidal anti-inflammatory agents. The variety of compounds encompassed by this group are well-known to those skilled in the art. For detailed disclosure of the chemical structure, synthesis, side effects, etc. of non-steroidal anti-inflammatory agents, one may refer to standard texts, including Anti-inflammatory and Anti-Rheumatic Drugs, K. D. Rainsford, Vol. I-III, CRC Press, Boca Raton, (1985), and Anti-inflammatory Agents, Chemistry and Pharmacology, 1, R. A. Scherrer, et al., Academic Press, New York (1974).

Specific non-steroidal anti-inflammatory agents useful in the composition invention include, but are not limited to:

- 1) the oxicams, such as piroxicam, isoxicam, tenoxicam, sudoxicam, and CP-14,304;
- 2) the salicylates, such as aspirin, disalcid, benorylate, trilisate, safapryn, solprin, diflunisal, and fendosal;
- 3) the acetic acid derivatives, such as diclofenac, fenclofenac, indomethacin, sulindac, tolmetin, isoxepac, furofenac, tiopinac, zidometacin, acematacin, fentiazac, zomepirac, clindanac, oxepinac, felbinac, and ketorolac;
- 4) the fenamates, such as mefenamic, meclofenamic, flufenamic, niflumic, and tolfenamic acids;

5) the propionic acid derivatives, such as ibuprofen, naproxen, benoxaprofen, flurbiprofen, ketoprofen, fenoprofen, fenbufen, indoprofen, pirofen, carprofen, oxaprozin, pranoprofen, miroprofen, tioxaprofen, suprofen, alminoprofen, and tiaprofenic; and

6) the pyrazoles, such as phenylbutazone, oxyphenbutazone, feprazone, azapropazone, and trimethazone.

Mixtures of these non-steroidal anti-inflammatory agents may also be employed, as well as the dermatologically acceptable salts and esters of these agents. For example, etofenamate, a flufenamic acid derivative, is particularly useful for topical application. Of the nonsteroidal anti-inflammatory agents, ibuprofen, naproxen, flufenamic acid, etofenamate, aspirin, mefenamic acid, meclofenamic acid, piroxicam and felbinac are preferred; ibuprofen, naproxen, ketoprofen, etofenamate, aspirin and flufenamic acid are more preferred.

Finally, so-called "natural" anti-inflammatory agents are useful in methods of the present invention. Such agents may suitably be obtained as an extract by suitable physical and/or chemical isolation from natural sources (e.g., plants, fungi, by-products of microorganisms) or can be synthetically prepared. For example, candelilla wax, bisabolol (e.g., alpha bisabolol), aloe vera, plant sterols (e.g., phytosterol), Manjistha (extracted from plants in the genus Rubia, particularly Rubia Cordifolia), and Guggal (extracted from plants in the genus Commiphora, particularly Commiphora Mukul), kola extract, chamomile, red clover extract, and sea whip extract, may be used.

Additional anti-inflammatory agents useful herein include compounds of the Licorice (the plant genus/species Glycyrrhiza glabra) family, including glycyrrhetic acid, glycyrrhizic acid, and derivatives thereof (e.g., salts and esters). Suitable salts of the foregoing compounds include metal and ammonium salts. Suitable esters include C₂ - C₂₄ saturated or unsaturated esters of the acids, preferably C₁₀ - C₂₄, more preferably C₁₆ - C₂₄. Specific examples of the foregoing include oil soluble licorice extract, the glycyrrhizic and glycyrrhetic acids themselves, monoammonium glycyrrhizinate, monopotassium glycyrrhizinate, dipotassium glycyrrhizinate, 1-beta-glycyrrhetic acid, stearyl glycyrrhetinate, and 3-stearyloxy-glycyrrhetinic acid, and disodium 3-succinyloxy-beta-glycyrrhetinate. Stearyl glycyrrhetinate is preferred.

Anti-Cellulite Agents

The compositions of the present invention may also contain a safe and effective amount of an anti-cellulite agent. Suitable agents may include, but are not limited to, xanthine compounds (e.g., caffeine, theophylline, theobromine, and aminophylline).


Topical Anesthetics

The compositions of the present invention may also contain a safe and effective amount of a topical anesthetic. Examples of topical anesthetic drugs include benzocaine, lidocaine, bupivacaine, chlorprocaine, dibucaine, etidocaine, mepivacaine, tetracaine, dyclonine, hexylcaine, procaine, cocaine, ketamine, pramoxine, phenol, and pharmaceutically acceptable salts thereof.

Tanning Actives

The compositions of the present invention may contain a tanning active. When present, it is preferable that the compositions contain from about 0.1% to about 20%, more preferably from about 2% to about 7%, and still more preferably from about 3% to about 6%, by weight of the composition, of dihydroxyacetone as an artificial tanning active.

Dihydroxyacetone, which is also known as DHA or 1,3-dihydroxy-2-propanone, is a white to off-white, crystalline powder. This material can be represented by the chemical formula $C_3H_6O_3$ and the following chemical structure.



The compound can exist as a mixture of monomers and dimers, with the dimers predominating in the solid crystalline state. Upon heating or melting, the dimers break down to yield the monomers. This conversion of the dimeric form to the monomeric form also occurs in aqueous solution. Dihydroxyacetone is also known to be more stable at acidic pH values. See The Merck Index, Tenth Edition, entry 3167, p. 463 (1983), and "Dihydroxyacetone for Cosmetics", E. Merck Technical Bulletin, 03-304 110, 319 897, 180 588.

Skin Lightening Agents

The compositions of the present invention may contain a skin lightening agent. When used, the compositions preferably contain from about 0.1% to about 10%, more preferably from about 0.2% to about 5%, also preferably from about 0.5% to about 2%, by weight of the composition, of a skin lightening agent. Suitable skin lightening agents include those known in the art, including kojic acid, arbutin, ascorbic acid and derivatives thereof (e.g., magnesium ascorbyl phosphate or sodium ascorbyl phosphate), and extracts (e.g., mulberry extract, placental extract). Skin lightening agents suitable for use herein also include those described in the PCT publication No. 95/34280, in the name of Hillebrand, corresponding to PCT

Application No. U.S. 95/07432, filed 6/12/95; and co-pending U.S. Application No. 08/390,152 filed in the names of Kvalnes, Mitchell A. DeLong, Barton J. Bradbury, Curtis B. Motley, and John D. Carter, corresponding to PCT Publication No. 95/23780, published 9/8/95.

Skin Soothing and Skin Healing Actives

The compositions of the present invention may contain a skin soothing or skin healing active. Skin soothing or skin healing actives suitable for use herein include panthenoic acid derivatives (including panthenol, dexpanthenol, ethyl panthenol), aloe vera, allantoin, bisabolol, and dipotassium glycyrrhizinate. A safe and effective amount of a skin soothing or skin healing active may be added to the present composition, preferably, from about 0.1% to about 30%, more preferably from about 0.5% to about 20%, still more preferably from about 0.5% to about 10 %, by weight of the composition formed.

Antimicrobial and Antifungal Actives

The compositions of the present invention may contain an antimicrobial or antifungal active. Such actives are capable of destroying microbes, preventing the development of microbes or preventing the pathogenic action of microbes. A safe and effective amount of an antimicrobial or antifungal active may be added to the present compositions, preferably, from about 0.001% to about 10%, more preferably from about 0.01% to about 5%, and still more preferably from about 0.05% to about 2%.

Examples of antimicrobial and antifungal actives include β -lactam drugs, quinolone drugs, ciprofloxacin, norfloxacin, tetracycline, erythromycin, amikacin, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorobanilide, phenoxyethanol, phenoxy propanol, phenoxyisopropanol, doxycycline, capreomycin, chlorhexidine, chlortetracycline, oxytetracycline, clindamycin, ethambutol, hexamidine isethionate, metronidazole, pentamidine, gentamicin, kanamycin, lineomycin, methacycline, methenamine, minocycline, neomycin, netilmicin, paromomycin, streptomycin, tobramycin, miconazole, tetracycline hydrochloride, erythromycin, zinc erythromycin, erythromycin estolate, erythromycin stearate, amikacin sulfate, doxycycline hydrochloride, capreomycin sulfate, chlorhexidine gluconate, chlorhexidine hydrochloride, chlortetracycline hydrochloride, oxytetracycline hydrochloride, clindamycin hydrochloride, ethambutol hydrochloride, metronidazole hydrochloride, pentamidine hydrochloride, gentamicin sulfate, kanamycin sulfate, lineomycin hydrochloride, methacycline hydrochloride, methenamine hippurate, methenamine mandelate, minocycline hydrochloride, neomycin sulfate, netilmicin sulfate, paromomycin sulfate, streptomycin sulfate, tobramycin

sulfate, miconazole hydrochloride, ketaconazole, amantadine hydrochloride, amantadine sulfate, octopirox, parachlorometa xylenol, nystatin, tolnaftate, zinc pyrithione and clotrimazole.

Preferred examples of actives useful herein include those selected from salicylic acid, benzoyl peroxide, 3-hydroxy benzoic acid, glycolic acid, lactic acid, 4-hydroxy benzoic acid, acetyl salicylic acid, 2-hydroxybutanoic acid, 2-hydroxypentanoic acid, 2-hydroxyhexanoic acid, cis-retinoic acid, trans-retinoic acid, retinol, phytic acid, N-acetyl-L-cysteine, lipoic acid, azelaic acid, arachidonic acid, benzoylperoxide, tetracycline, ibuprofen, naproxen, hydrocortisone, acetaminophen, resorcinol, phenoxyethanol, phenoxypropanol, phenoxyisopropanol, 2,4,4'-trichloro-2'-hydroxy diphenyl ether, 3,4,4'-trichlorocarbaniide, octopirox, lidocaine hydrochloride, clotrimazole, miconazole, ketoconazole, neocycin sulfate, and mixtures thereof.

Sunscreen Actives

Exposure to ultraviolet light can result in excessive scaling and texture changes of the stratum corneum. Therefore, the compositions of the subject invention may optionally contain a sunscreen active. As used herein, "sunscreen active" includes both sunscreen agents and physical sunblocks. Suitable sunscreen actives may be organic or inorganic.

Inorganic sunscreens useful herein include the following metallic oxides; titanium dioxide having an average primary particle size of from about 15 nm to about 100 nm, zinc oxide having an average primary particle size of from about 15 nm to about 150 nm, zirconium oxide having an average primary particle size of from about 15 nm to about 150 nm, iron oxide having an average primary particle size of from about 15 nm to about 500nm, and mixtures thereof. When used herein, the inorganic sunscreens are present in the amount of from about 0.1% to about 20%, preferably from about 0.5% to about 10%, more preferably from about 1% to about 5%, by weight of the composition.

A wide variety of conventional organic sunscreen actives are suitable for use herein. Sagarin, et al., at Chapter VIII, pages 189 et seq., of Cosmetics Science and Technology (1972), discloses numerous suitable actives. Specific suitable sunscreen actives include, for example: p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-dimethylaminobenzoic acid); anthranilates (i.e., o-amino-benzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (amyl, phenyl, octyl, benzyl, menthyl, glyceryl, and di-pro-pyleneglycol esters); cinnamic acid derivatives (menthyl and benzyl esters, a-phenyl cinnamionitrile; butyl cinnamoyl pyruvate); dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); trihydroxy-

cinnamic acid derivatives (esculetin, methylesculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone and benzalacetophenone; naphtholsulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); di-hydroxynaphthoic acid and its salts; o- and p-hydroxybiphenyldisulfonates; coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); diazoles (2-acetyl-3-bromoindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); hydroxy- or methoxy-substituted benzophenones; uric and violuric acids; tannic acid and its derivatives (e.g., hexaethylether); (butyl carbotol) (6-propyl piperonyl) ether; hydroquinone; benzophenones (oxybenzene, sulisobenzene, dioxybenzene, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-dihydroxy-4,4'-dimethoxybenzophenone, octabenzene; 4-isopropylidibenzoylmethane; butylmethoxydibenzoylmethane; etocrylene; octocrylene; [3-(4'-methylbenzylidene bornan-2-one), terephthalylidene dicamphor sulfonic acid and 4-isopropyl-di-benzoylmethane.

Of these, 2-ethylhexyl-p-methoxycinnamate (commercially available as PARSOL MCX), 4,4'-t-butyl methoxydibenzoyl-methane (commercially available as PARSOL 1789), 2-hydroxy-4-methoxybenzophenone, octyldimethyl-p-aminobenzoic acid, digalloyltrioleate, 2,2'-dihydroxy-4-methoxybenzophenone, ethyl-4-(bis(hydroxy-propyl))aminobenzoate, 2-ethylhexyl-2-cyano-3,3-diphenylacrylate, 2-ethylhexyl-salicylate, glyceryl-p-aminobenzoate, 3,3,5-tri-methylcyclohexylsalicylate, methylanthranilate, p-dimethyl-aminobenzoic acid or aminobenzoate, 2-ethylhexyl-p-dimethyl-amino-benzoate, 2-phenylbenzimidazole-5-sulfonic acid, 2-(p-dimethylaminophenyl)-5-sulfonicbenzoxazoic acid, octocrylene and mixtures of these compounds, are preferred.

More preferred organic sunscreen actives useful in the compositions useful in the subject invention are 2-ethylhexyl-p-methoxycinnamate, butylmethoxydibenzoyl-methane, 2-hydroxy-4-methoxybenzo-phenone, 2-phenylbenzimidazole-5-sulfonic acid, octyldimethyl-p-aminobenzoic acid, octocrylene and mixtures thereof.

Also particularly useful in the compositions are sunscreen actives such as those disclosed in U.S. Patent No. 4,937,370 issued to Sabatelli on June 26, 1990, and U.S. Patent No. 4,999,186 issued to Sabatelli & Spirnak on March 12, 1991. The sunscreens disclosed therein have, in a single molecule, two distinct chromophore moieties which exhibit different ultra-violet radiation absorption spectra. One of the chromophore moieties absorbs

predominantly in the UVB radiation range and the other absorbs strongly in the UVA radiation range.

Preferred members of this class of sunscreens are 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2,4-dihydroxybenzophenone; N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester with 4-hydroxydibenzoylmethane; 4-N,N-(2-ethylhexyl)methylaminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone; 4-N,N-(2-ethylhexyl)-methylaminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane; N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 2-hydroxy-4-(2-hydroxyethoxy)benzophenone; and N,N-di-(2-ethylhexyl)-4-aminobenzoic acid ester of 4-(2-hydroxyethoxy)dibenzoylmethane and mixtures thereof.

Especially preferred sunscreen actives include 4,4'-t-butylmethoxydibenzoylmethane, 2-ethylhexyl-p-methoxycinnamate, phenyl benzimidazole sulfonic acid, and octocrylene.

A safe and effective amount of the organic sunscreen active is used, typically from about 1% to about 20%, more typically from about 2% to about 10% by weight of the composition. Exact amounts will vary depending upon the sunscreen or sunscreens chosen and the desired Sun Protection Factor (SPF).

Particulate Material

The compositions of the present invention may contain a particulate material, preferably a metallic oxide. These particulates can be coated or uncoated, charged or uncharged. Charged particulate materials are disclosed in U.S. Patent No. 5,997,887, to Ha, et al., incorporated herein by reference. Particulate materials useful herein include; bismuth oxychloride, iron oxide, mica, mica treated with barium sulfate and TiO₂, silica, nylon, polyethylene, talc, styrene, polypropylene, ethylene/acrylic acid copolymer, titanium dioxide, iron oxide, bismuth oxychloride, sericite, aluminum oxide, silicone resin, barium sulfate, calcium carbonate, cellulose acetate, polymethyl methacrylate, and mixtures thereof.

Inorganic particulate materials, e.g., TiO₂, ZnO, or ZrO₂ are commercially available from a number of sources. One example of a suitable particulate material contains the material available from U.S. Cosmetics (TRONOX TiO₂ series, SAT-T CR837, a rutile TiO₂). Preferably, particulate materials are present in the composition in levels of from about 0.01% to about 2%, more preferably from about 0.05% to about 1.5%, still more preferably from about 0.1% to about 1%, by weight of the composition.

Conditioning Agents

The compositions of the present invention may contain a conditioning agent selected from humectants, moisturizers, or skin conditioners. A variety of these materials can be employed and each can be present at a level of from about 0.01% to about 20%, more preferably from about 0.1% to about 10%, and still more preferably from about 0.5% to about 7% by weight of the composition. These materials include, but are not limited to, guanidine; urea; glycolic acid and glycolate salts (e.g. ammonium and quaternary alkyl ammonium); salicylic acid; lactic acid and lactate salts (e.g., ammonium and quaternary alkyl ammonium); aloe vera in any of its variety of forms (e.g., aloe vera gel); polyhydroxy alcohols such as sorbitol, mannitol, xylitol, erythritol, glycerol, hexanetriol, butanetriol, propylene glycol, butylene glycol, hexylene glycol and the like; polyethylene glycols; sugars (e.g., melibiose) and starches; sugar and starch derivatives (e.g., alkoxylated glucose, fucose, glucosamine); hyaluronic acid; lactamide monoethanolamine; acetamide monoethanolamine; panthenol; allantoin; and mixtures thereof. Also useful herein are the propoxylated glycerols described in U. S. Patent No. 4,976,953, to Orr et al, issued December 11, 1990.

Also useful are various C₁-C₃₀ monoesters and polyesters of sugars and related materials. These esters are derived from a sugar or polyol moiety and one or more carboxylic acid moieties. Such ester materials are further described in, U. S. Patent No. 2,831,854, U. S. Patent No. 4,005,196, to Jandacek, issued January 25, 1977; U. S. Patent No. 4,005,195, to Jandacek, issued January 25, 1977, U. S. Patent No. 5,306,516, to Letton et al, issued April 26, 1994; U. S. Patent No. 5,306,515, to Letton et al, issued April 26, 1994; U. S. Patent No. 5,305,514, to Letton et al, issued April 26, 1994; U. S. Patent No. 4,797,300, to Jandacek et al, issued January 10, 1989; U. S. Patent No. 3,963,699, to Rizzi et al, issued June 15, 1976; U. S. Patent No. 4,518,772, to Volpenhein, issued May 21, 1985; and U. S. Patent No. 4,517,360, to Volpenhein, issued May 21, 1985.

Preferably, the conditioning agent is selected from urea, guanidine, sucrose polyester, panthenol, dexpanthenol, allantoin, and combinations thereof.

Structuring Agents

The compositions hereof, and especially the emulsions hereof, may contain a structuring agent. Structuring agents are particularly preferred in the oil-in-water emulsions of the present invention. Without being limited by theory, it is believed that the structuring agent assists in providing rheological characteristics to the composition which contribute to the stability of the composition. For example, the structuring agent tends to assist in the formation of the liquid

crystalline gel network structures. The structuring agent may also function as an emulsifier or surfactant. Preferred compositions of this invention contain from about 0.1% to about 20%, more preferably from about 0.1% to about 10%, still more preferably from about 0.5% to about 9%, of one or more structuring agents.

Preferred structuring agents are those having an HLB of from about 1 to about 8 and having a melting point of at least about 45°C. Suitable structuring agents are those selected from saturated C₁₄ to C₃₀ fatty alcohols, saturated C₁₆ to C₃₀ fatty alcohols containing from about 1 to about 5 moles of ethylene oxide, saturated C₁₆ to C₃₀ diols, saturated C₁₆ to C₃₀ monoglycerol ethers, saturated C₁₆ to C₃₀ hydroxy fatty acids, C₁₄ to C₃₀ hydroxylated and nonhydroxylated saturated fatty acids, C₁₄ to C₃₀ saturated ethoxylated fatty acids, amines and alcohols containing from about 1 to about 5 moles of ethylene oxide diols, C₁₄ to C₃₀ saturated glyceryl mono esters with a monoglyceride content of at least 40%, C₁₄ to C₃₀ saturated polyglycerol esters having from about 1 to about 3 alkyl group and from about 2 to about 3 saturated glycerol units, C₁₄ to C₃₀ glyceryl mono ethers, C₁₄ to C₃₀ sorbitan mono/diesters, C₁₄ to C₃₀ saturated ethoxylated sorbitan mono/diesters with about 1 to about 5 moles of ethylene oxide, C₁₄ to C₃₀ saturated methyl glucoside esters, C₁₄ to C₃₀ saturated sucrose mono/diesters, C₁₄ to C₃₀ saturated ethoxylated methyl glucoside esters with about 1 to about 5 moles of ethylene oxide, C₁₄ to C₃₀ saturated polyglucosides having an average of between 1 to 2 glucose units and mixtures thereof, having a melting point of at least about 45°C.

The preferred structuring agents of the present invention are selected from stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, stearic acid, palmitic acid, the polyethylene glycol ether of stearyl alcohol having an average of about 1 to about 5 ethylene oxide units, the polyethylene glycol ether of cetyl alcohol having an average of about 1 to about 5 ethylene oxide units, and mixtures thereof. More preferred structuring agents of the present invention are selected from stearyl alcohol, cetyl alcohol, behenyl alcohol, the polyethylene glycol ether of stearyl alcohol having an average of about 2 ethylene oxide units (steareth-2), the polyethylene glycol ether of cetyl alcohol having an average of about 2 ethylene oxide units, and mixtures thereof. Even more preferred structuring agents are selected from stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, steareth-2, and mixtures thereof.

Thickening Agent (including thickeners and gelling agents)

The compositions of the present invention can contain one or more thickening agents, preferably from about 0.1% to about 5%, more preferably from about 0.1% to about 4%, and still more preferably from about 0.25% to about 3%, by weight of the composition.

Nonlimiting classes of thickening agents include those selected from the following:

a) Carboxylic Acid Polymers

These polymers are crosslinked compounds containing one or more monomers derived from acrylic acid, substituted acrylic acids, and salts and esters of these acrylic acids and the substituted acrylic acids, wherein the crosslinking agent contains two or more carbon-carbon double bonds and is derived from a polyhydric alcohol. Polymers useful in the present invention are more fully described in U. S. Patent No. 5,087,445, to Haffey et al, issued February 11, 1992; U. S. Patent No. 4,509,949, to Huang et al, issued April 5, 1985; U. S. Patent No. 2,798,053, to Brown, issued July 2, 1957; and in *CTFA International Cosmetic Ingredient Dictionary*, Fourth Edition, 1991, pp. 12 and 80.

Examples of commercially available carboxylic acid polymers useful herein include the carbomers, which are homopolymers of acrylic acid crosslinked with allyl ethers of sucrose or pentaerythritol. The carbomers are available as the Carbopol® 900 series from B.F. Goodrich (e.g., Carbopol® 954). In addition, other suitable carboxylic acid polymeric agents include copolymers of C₁₀₋₃₀ alkyl acrylates with one or more monomers of acrylic acid, methacrylic acid, or one of their short chain (i.e., C₁₋₄ alcohol) esters, wherein the crosslinking agent is an allyl ether of sucrose or pentaerythritol. These copolymers are known as acrylates/C₁₀₋₃₀ alkyl acrylate crosspolymers and are commercially available as Carbopol® 1342, Carbopol® 1382, Pemulen TR-1, and Pemulen TR-2, from B.F. Goodrich. In other words, examples of carboxylic acid polymer thickeners useful herein are those selected from carbomers, acrylates/C_{10-C₃₀} alkyl acrylate crosspolymers, and mixtures thereof.

b) Crosslinked Polyacrylate Polymers

The compositions of the present invention can optionally contain crosslinked polyacrylate polymers useful as thickeners or gelling agents including both cationic and nonionic polymers, with the cationics being generally preferred. Examples of useful crosslinked nonionic polyacrylate polymers and crosslinked cationic polyacrylate polymers are those described in U. S. Patent No. 5,100,660, to Hawe et al, issued March 31, 1992; U. S. Patent No.

4,849,484, to Heard, issued July 18, 1989; U. S. Patent No. 4,835,206, to Farrar et al, issued May 30, 1989; U.S. Patent No. 4,628,078 to Glover et al issued December 9, 1986; U.S. Patent No. 4,599,379 to Flesher et al issued July 8, 1986; and EP 228,868, to Farrar et al, published July 15, 1987.

c) Polyacrylamide Polymers

The compositions of the present invention can optionally contain polyacrylamide polymers, especially nonionic polyacrylamide polymers including substituted branched or unbranched polymers. More preferred among these polyacrylamide polymers is the nonionic polymer given the CTFA designation polyacrylamide and isoparaffin and laureth-7, available under the Tradename Sepigel 305 from Seppic Corporation (Fairfield, NJ).

Other polyacrylamide polymers useful herein include multi-block copolymers of acrylamides and substituted acrylamides with acrylic acids and substituted acrylic acids. Commercially available examples of these multi-block copolymers include Hypan SR150H, SS500V, SS500W, SSSA100H, from Lipo Chemicals, Inc., (Patterson, NJ).

d) Polysaccharides

A wide variety of polysaccharides are useful herein. "Polysaccharides" refer to gelling agents which contain a backbone of repeating sugar (i.e., carbohydrate) units. Nonlimiting examples of polysaccharide gelling agents include those selected from cellulose, carboxymethyl hydroxyethylcellulose, cellulose acetate propionate carboxylate, hydroxyethylcellulose, hydroxyethyl ethylcellulose, hydroxypropylcellulose, hydroxypropyl methylcellulose, methyl hydroxyethylcellulose, microcrystalline cellulose, sodium cellulose sulfate, and mixtures thereof. Also useful herein are the alkyl substituted celluloses. In these polymers, the hydroxy groups of the cellulose polymer is hydroxyalkylated (preferably hydroxyethylated or hydroxypropylated) to form a hydroxyalkylated cellulose which is then further modified with a C₁₀-C₃₀ straight chain or branched chain alkyl group through an ether linkage. Typically these polymers are ethers of C₁₀-C₃₀ straight or branched chain alcohols with hydroxyalkylcelluloses. Examples of alkyl groups useful herein include those selected from stearyl, isostearyl, lauryl, myristyl, cetyl, isocetyl, cocoyl (i.e. alkyl groups derived from the alcohols of coconut oil), palmityl, oleyl, linoleyl, linolenyl, ricinoleyl, behenyl, and mixtures thereof. Preferred among the alkyl hydroxyalkyl cellulose ethers is the material given the CTFA designation cetyl hydroxyethylcellulose, which is the ether of cetyl alcohol and hydroxyethylcellulose. This material is sold under the tradename Natrosol® CS Plus from Aqualon Corporation (Wilmington, DE).

Other useful polysaccharides include scleroglucans which are a linear chain of (1-3) linked glucose units with a (1-6) linked glucose every three units, a commercially available example of which is Clearogel™ CS11 from Michel Mercier Products Inc. (Mountainside, NJ).

e) Gums

Other thickening and gelling agents useful herein include materials which are primarily derived from natural sources. Nonlimiting examples of these gelling agent gums include acacia, agar, algin, alginic acid, ammonium alginate, amylopectin, calcium alginate, calcium carrageenan, carnitine, carrageenan, dextrin, gelatin, gellan gum, guar gum, guar hydroxypropyltrimonium chloride, hectorite, hyaluronic acid, hydrated silica, hydroxypropyl chitosan, hydroxypropyl guar, karaya gum, kelp, locust bean gum, natto gum, potassium alginate, potassium carrageenan, propylene glycol alginate, sclerotium gum, sodium carboxymethyl dextran, sodium carrageenan, tragacanth gum, xanthan gum, and mixtures thereof.

Preferred compositions of the present invention include a thickening agent selected from carboxylic acid polymers, crosslinked polyacrylate polymers, polyacrylamide polymers, and mixtures thereof, more preferably selected from carboxylic acid polymers, polyacrylamide polymers, and mixtures thereof.

IV. Dermatologically-Acceptable Carrier

The topical compositions of the present invention also contain a dermatologically acceptable carrier. The phrase "dermatologically-acceptable carrier", as used herein, means that the carrier is suitable for topical application to the keratinous tissue, has good aesthetic properties, is compatible with the actives of the present invention and any other components, and will not cause any untoward safety or toxicity concerns. A safe and effective amount of carrier is from about 50% to about 99.99%, preferably from about 80% to about 99.9%, more preferably from about 90% to about 98%, and even more preferably from about 90% to about 95% of the composition.

The carrier can be in a wide variety of forms. For example, emulsion carriers, including, but not limited to, oil-in-water, water-in-oil, water-in-oil-in-water, and oil-in-water-in-silicone emulsions, are useful herein.

Preferred carriers contain an emulsion such as oil-in-water emulsions, water-in-oil emulsions, and water-in-silicone emulsions. As will be understood by the skilled artisan, a given component will distribute primarily into either the water or oil/silicone phase, depending

on the water solubility/dispersibility of the component in the composition. Oil-in-water emulsions are especially preferred.

Emulsions according to the present invention generally contain a solution as described above and a lipid or oil. Lipids and oils may be derived from animals, plants, or petroleum and may be natural or synthetic (i.e., man-made). Preferred emulsions also contain a humectant, such as glycerin. Emulsions will preferably further contain from about 0.01% to about 10%, more preferably from about 0.1% to about 5%, of an emulsifier, based on the weight of the carrier. Emulsifiers may be nonionic, anionic or cationic. Suitable emulsifiers are disclosed in, for example, U.S. Patent 3,755,560, issued August 28, 1973, Dickert et al.; U.S. Patent 4,421,769, issued December 20, 1983, Dixon et al.; and McCutcheon's Detergents and Emulsifiers, North American Edition, pages 317-324 (1986).

The emulsion may also contain an anti-foaming agent to minimize foaming upon application to the keratinous tissue. Anti-foaming agents include high molecular weight silicones and other materials well known in the art for such use.

Suitable emulsions may have a wide range of viscosities, depending on the desired product form. Exemplary low viscosity emulsions, which are preferred, have a viscosity of about 50 centistokes or less, more preferably about 10 centistokes or less, still more preferably about 5 centistokes or less.

Preferred water-in-silicone and oil-in-water emulsions are described in greater detail below.

A) Water-in-silicone emulsion

Water-in-silicone emulsions contain a continuous silicone phase and a dispersed aqueous phase.

(1) Continuous silicone phase

Preferred water-in-silicone emulsions of the present invention contain from about 1% to about 60%, preferably from about 5% to about 40%, more preferably from about 10% to about 20%, by weight of a continuous silicone phase. The continuous silicone phase exists as an external phase that contains or surrounds the discontinuous aqueous phase described hereinafter.

The continuous silicone phase contains a polyorganosiloxane oil. A preferred water-in-silicone emulsion system is formulated to provide an oxidatively stable vehicle for the retinoid. The continuous silicone phase of these preferred emulsions contain between about 50% and about 99.9% by weight of organopolysiloxane oil and less than about 50% by weight of a non-silicone oil. In an especially preferred embodiment, the continuous silicone phase contains at

least about 50%, preferably from about 60% to about 99.9%, more preferably from about 70% to about 99.9%, and even more preferably from about 80% to about 99.9%, polyorganosiloxane oil by weight of the continuous silicone phase, and up to about 50% non-silicone oils, preferably less about 40%, more preferably less than about 30%, even more preferably less than about 10%, and even more preferably less than about 2%, by weight of the continuous silicone phase. These preferred emulsion systems provide more oxidative stability to the retinoid over extended periods of time than comparable water-in-oil emulsions containing lower concentrations of the polyorganosiloxane oil. Concentrations of non-silicone oils in the continuous silicone phase are minimized or avoided altogether so as to further enhance oxidative stability of the selected retinoid in the compositions. Water-in-silicone emulsions of this type are described in PCT Application WO 97/21423, published June 19, 1997.

The organopolysiloxane oil for use in the composition may be volatile, non-volatile, or a mixture of volatile and non-volatile silicones. The term "nonvolatile" as used in this context refers to those silicones that are liquid under ambient conditions and have a flash point (under one atmospheric of pressure) of or greater than about 100°C. The term "volatile" as used in this context refers to all other silicone oils. Suitable organopolysiloxanes can be selected from a wide variety of silicones spanning a broad range of volatilities and viscosities. Examples of suitable organopolysiloxane oils include polyalkylsiloxanes, cyclic polyalkylsiloxanes, and polyalkylarylsiloxanes.

Polyalkylsiloxanes useful in the composition herein include polyalkylsiloxanes with viscosities of from about 0.5 to about 1,000,000 centistokes at 25°C. Such polyalkylsiloxanes can be represented by the general chemical formula $R_3SiO[R_2SiO]_xSiR_3$ wherein R is an alkyl group having from one to about 30 carbon atoms (preferably R is methyl or ethyl, more preferably methyl; also mixed alkyl groups can be used in the same molecule), and x is an integer from 0 to about 10,000, chosen to achieve the desired molecular weight which can range to over about 10,000,000. Commercially available polyalkylsiloxanes include the polydimethylsiloxanes, which are also known as dimethicones, examples of which include the Vicasil[®] series sold by General Electric Company and the Dow Corning[®] 200 series sold by Dow Corning Corporation. Specific examples of suitable polydimethylsiloxanes include Dow Corning[®] 200 fluid having a viscosity of 0.65 centistokes and a boiling point of 100°C, Dow Corning[®] 225 fluid having a viscosity of 10 centistokes and a boiling point greater than 200°C, and Dow Corning[®] 200 fluids having viscosities of 50, 350, and 12,500 centistokes,

respectively, and boiling points greater than 200°C. Suitable dimethicones include those represented by the chemical formula $(\text{CH}_3)_3\text{SiO}[(\text{CH}_3)_2\text{SiO}]_x[\text{CH}_3\text{RSiO}]_y\text{Si}(\text{CH}_3)_3$ wherein R is straight or branched chain alkyl having from two to about 30 carbon atoms and x and y are each integers of 1 or greater selected to achieve the desired molecular weight which can range to over about 10,000,000. Examples of these alkyl-substituted dimethicones include cetyl dimethicone and lauryl dimethicone.

Cyclic polyalkylsiloxanes suitable for use in the composition include those represented by the chemical formula $[\text{SiR}_2\text{-O}]_n$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and n is an integer from about 3 to about 8, more preferably n is an integer from about 3 to about 7, and still more preferably n is an integer from about 4 to about 6. When R is methyl, these materials are typically referred to as cyclomethicones. Commercially available cyclomethicones include Dow Corning® 244 fluid having a viscosity of 2.5 centistokes, and a boiling point of 172°C, which primarily contains the cyclomethicone tetramer (i.e. n=4), Dow Corning® 344 fluid having a viscosity of 2.5 centistokes and a boiling point of 178°C, which primarily contains the cyclomethicone pentamer (i.e. n=5), Dow Corning® 245 fluid having a viscosity of 4.2 centistokes and a boiling point of 205°C, which primarily contains a mixture of the cyclomethicone tetramer and pentamer (i.e. n=4 and 5), and Dow Corning® 345 fluid having a viscosity of 4.5 centistokes and a boiling point of 217°, which primarily contains a mixture of the cyclomethicone tetramer, pentamer, and hexamer (i.e. n=4, 5, and 6).

Also useful are materials such as trimethylsiloxysilicate, which is a polymeric material corresponding to the general chemical formula $[(\text{CH}_3)_3\text{SiO}_{1/2}]_x[\text{SiO}_2]_y$, wherein x is an integer from about 1 to about 500 and y is an integer from about 1 to about 500. A commercially available trimethylsiloxysilicate is sold as a mixture with dimethicone as Dow Corning® 593 fluid.

Dimethiconols are also suitable for use in the composition. These compounds can be represented by the chemical formulas $\text{R}_3\text{SiO}[\text{R}_2\text{SiO}]_x\text{SiR}_2\text{OH}$ and $\text{HOR}_2\text{SiO}[\text{R}_2\text{SiO}]_x\text{SiR}_2\text{OH}$ wherein R is an alkyl group (preferably R is methyl or ethyl, more preferably methyl) and x is an integer from 0 to about 500, chosen to achieve the desired molecular weight. Commercially available dimethiconols are typically sold as mixtures with dimethicone or cyclomethicone (e.g. Dow Corning® 1401, 1402, and 1403 fluids).

Polyalkylaryl siloxanes are also suitable for use in the composition. Polymethylphenyl siloxanes having viscosities from about 15 to about 65 centistokes at 25°C are especially useful.

Preferred for use herein are organopolysiloxanes selected from polyalkylsiloxanes, alkyl substituted dimethicones, cyclomethicones, trimethylsiloxysilicates, dimethiconols, polyalkylaryl siloxanes, and mixtures thereof. More preferred for use herein are polyalkylsiloxanes and cyclomethicones. Preferred among the polyalkylsiloxanes are dimethicones.

As stated above, the continuous silicone phase may contain one or more non-silicone oils. Concentrations of non-silicone oils in the continuous silicone phase are preferably minimized or avoided altogether so as to further enhance oxidative stability of the selected retinoid in the compositions. Suitable non-silicone oils have a melting point of about 25°C or less under about one atmosphere of pressure. Examples of non-silicone oils suitable for use in the continuous silicone phase are those well known in the chemical arts in topical personal care products in the form of water-in-oil emulsions, e.g., mineral oil, vegetable oils, synthetic oils, semisynthetic oils, etc.

(2) Dispersed aqueous phase

The topical compositions of the present invention contain from about 30% to about 90%, more preferably from about 50% to about 85%, and still more preferably from about 70% to about 80% of a dispersed aqueous phase. In emulsion technology, the term "dispersed phase" is a term well-known to one skilled in the art which means that the phase exists as small particles or droplets that are suspended in and surrounded by a continuous phase. The dispersed phase is also known as the internal or discontinuous phase. The dispersed aqueous phase is a dispersion of small aqueous particles or droplets suspended in and surrounded by the continuous silicone phase described hereinbefore.

The aqueous phase can be water, or a combination of water and one or more water soluble or dispersible ingredients. Nonlimiting examples of such ingredients include thickeners, acids, bases, salts, chelants, gums, water-soluble or dispersible alcohols and polyols, buffers, preservatives, sunscreens agents, colorings, and the like.

The topical compositions of the present invention will typically contain from about 25% to about 90%, preferably from about 40% to about 80%, more preferably from about 60% to about 80%, water in the dispersed aqueous phase by weight of the composition.


(3) Emulsifier for dispersing the aqueous phase

The water-in-silicone emulsions of the present invention preferably contain an emulsifier. In a preferred embodiment, the composition contains from about 0.1% to about 10% emulsifier, more preferably from about 0.5% to about 7.5%, still more preferably from about 1% to about 5%, emulsifier by weight of the composition. The emulsifier helps disperse and suspend the aqueous phase within the continuous silicone phase.

A wide variety of emulsifying agents can be employed herein to form the preferred water-in-silicone emulsion. Known or conventional emulsifying agents can be used in the composition, provided that the selected emulsifying agent is chemically and physically compatible with components of the composition of the present invention, and provides the desired dispersion characteristics. Suitable emulsifiers include silicone emulsifiers, non-silicon-containing emulsifiers, and mixtures thereof, known by those skilled in the art for use in topical personal care products. Preferably these emulsifiers have an HLB value of or less than about 14, more preferably from about 2 to about 14, and still more preferably from about 4 to about 14. Emulsifiers having an HLB value outside of these ranges can be used in combination with other emulsifiers to achieve an effective weighted average HLB for the combination that falls within these ranges.

Silicone emulsifiers are preferred. A wide variety of silicone emulsifiers are useful herein. These silicone emulsifiers are typically organically modified organopolysiloxanes, also known to those skilled in the art as silicone surfactants. Useful silicone emulsifiers include dimethicone copolyols. These materials are polydimethyl siloxanes which have been modified to include polyether side chains such as polyethylene oxide chains, polypropylene oxide chains, mixtures of these chains, and polyether chains containing moieties derived from both ethylene oxide and propylene oxide. Other examples include alkyl-modified dimethicone copolyols, i.e., compounds which contain C2-C30 pendant side chains. Still other useful dimethicone copolyols include materials having various cationic, anionic, amphoteric, and zwitterionic pendant moieties.


The dimethicone copolyol emulsifiers useful herein can be described by the following general structure:



wherein R is C1-C30 straight, branched, or cyclic alkyl and R² is selected from the group consisting of




and



wherein n is an integer from 3 to about 10; R³ and R⁴ are selected from the group consisting of H and C1-C6 straight or branched chain alkyl such that R³ and R⁴ are not simultaneously the same; and m, o, x, and y are selected such that the molecule has an overall molecular weight from about 200 to about 10,000,000, with m, o, x, and y being independently selected from integers of zero or greater such that m and o are not both simultaneously zero, and z being independently selected from integers of 1 or greater. It is recognized that positional isomers of these copolyols can be achieved. The chemical representations depicted above for the R² moieties containing the R³ and R⁴ groups are not meant to be limiting but are shown as such for convenience.

Also useful herein, although not strictly classified as dimethicone copolyols, are silicone surfactants as depicted in the structures in the previous paragraph wherein R² is:



wherein R⁵ is a cationic, anionic, amphoteric, or zwitterionic moiety.

Nonlimiting examples of dimethicone copolyols and other silicone surfactants useful as emulsifiers herein include polydimethylsiloxane polyether copolymers with pendant polyethylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant polypropylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant mixed polyethylene oxide and polypropylene oxide sidechains, polydimethylsiloxane polyether copolymers with pendant mixed poly(ethylene)(propylene)oxide sidechains, polydimethylsiloxane polyether copolymers with pendant organobetaine sidechains, polydimethylsiloxane polyether copolymers with pendant carboxylate sidechains, polydimethylsiloxane polyether copolymers with pendant quaternary ammonium sidechains; and also further modifications of the preceding copolymers containing pendant C2-C30 straight, branched, or cyclic alkyl moieties. Examples of commercially available dimethicone copolyols useful herein sold by Dow Corning Corporation are Dow Corning® 190, 193, Q2-5220, 2501 Wax, 2-5324 fluid, and 3225C (this later material being sold as a mixture with cyclomethicone).

Cetyl dimethicone copolyol is commercially available as a mixture with polyglyceryl-4 isostearate (and) hexyl laurate and is sold under the tradename ABIL[®] WE-09 (available from Goldschmidt). Cetyl dimethicone copolyol is also commercially available as a mixture with hexyl laurate (and) polyglyceryl-3 oleate (and) cetyl dimethicone and is sold under the tradename ABIL[®] WS-08 (also available from Goldschmidt). Other nonlimiting examples of dimethicone copolyols also include lauryl dimethicone copolyol, dimethicone copolyol acetate, diemethicone copolyol adipate, dimethicone copolyolamine, dimethicone copolyol behenate, dimethicone copolyol butyl ether, dimethicone copolyol hydroxy stearate, dimethicone copolyol isostearate, dimethicone copolyol laurate, dimethicone copolyol methyl ether, dimethicone copolyol phosphate, and dimethicone copolyol stearate. See International Cosmetic Ingredient Dictionary, Fifth Edition, 1993.

Dimethicone copolyol emulsifiers useful herein are described, for example, in U.S. Patent No. 4,960,764, to Figueroa, Jr. et al., issued October 2, 1990; European Patent No. EP 330,369, to SanoGueira, published August 30, 1989; G.H. Dahms, et al., "New Formulation Possibilities Offered by Silicone Copolyols," Cosmetics & Toiletries, vol. 110, pp. 91-100, March 1995; M.E. Carlotti et al., "Optimization of W/O-S Emulsions And Study Of The Quantitative Relationships Between Ester Structure And Emulsion Properties," J. Dispersion Science And Technology, 13(3), 315-336 (1992); P. Hameyer, "Comparative Technological Investigations of Organic and Organosilicone Emulsifiers in Cosmetic Water-in-Oil Emulsion Preparations," HAPPI 28(4), pp. 88-128 (1991); J. Smid-Korbar et al., "Efficiency and usability of silicone surfactants in emulsions," Provisional Communication, International Journal of Cosmetic Science, 12, 135-139 (1990); and D.G. Krzysik et al., "A New Silicone Emulsifier For Water-in-Oil Systems," Drug and Cosmetic Industry, vol. 146(4) pp. 28-81 (April 1990).

Among the non-silicone-containing emulsifiers useful herein are various non-ionic and anionic emulsifying agents such as sugar esters and polyesters, alkoxylated sugar esters and polyesters, C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxylated derivatives of C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxylated ethers of C1-C30 fatty alcohols, polyglyceryl esters of C1-C30 fatty acids, C1-C30 esters of polyols, C1-C30 ethers of polyols, alkyl phosphates, polyoxyalkylene fatty ether phosphates, fatty acid amides, acyl lactylates, soaps, and mixtures thereof. Other suitable emulsifiers are described, for example, in McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681 to Ciotti et al., issued April 30, 1991;

U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973.

Nonlimiting examples of these non-silicon-containing emulsifiers include: polyethylene glycol 20 sorbitan monolaurate (Polysorbate 20), polyethylene glycol 5 soya sterol, Steareth-20, Cetareth-20, PPG-2 methyl glucose ether distearate, Ceteth-10, Polysorbate 80, cetyl phosphate, potassium cetyl phosphate, diethanolamine cetyl phosphate, Polysorbate 60, glyceryl stearate, PEG-100 stearate, polyoxyethylene 20 sorbitan trioleate (Polysorbate 85), sorbitan monolaurate, polyoxyethylene 4 lauryl ether sodium stearate, polyglyceryl-4 isostearate, hexyl laurate, steareth-20, cetareth-20, PPG-2 methyl glucose ether distearate, ceteth-10, diethanolamine cetyl phosphate, glyceryl stearate, PEG-100 stearate, and mixtures thereof.

B) Oil-in-Water Emulsions

Other preferred topical carriers include oil-in-water emulsions, having a continuous aqueous phase and a hydrophobic, water-insoluble phase ("oil phase") dispersed therein. Examples of suitable oil-in-water emulsion carriers are described in U.S. Pat. No. 5,073,371, to Turner, D.J. et al., issued Dec. 17, 1991, and U.S. Pat. No. 5,073,372, to Turner, D.J. et al., issued Dec. 17, 1991. An especially preferred oil-in-water emulsion, containing a structuring agent, hydrophilic surfactant and water, is described in detail hereinafter.

(1) Structuring Agent

A preferred oil-in-water emulsion contains a structuring agent to assist in the formation of a liquid crystalline gel network structure. Without being limited by theory, it is believed that the structuring agent assists in providing rheological characteristics to the composition which contribute to the stability of the composition. The structuring agent may also function as an emulsifier or surfactant. Preferred compositions of this invention contain from about 0.5% to about 20%, more preferably from about 1% to about 10%, even more preferably from about 1% to about 5%, by weight of the composition, of a structuring agent.

The preferred structuring agents of the present invention include stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, stearic acid, palmitic acid, the polyethylene glycol ether of stearyl alcohol having an average of about 1 to about 21 ethylene oxide units, the polyethylene glycol ether of cetyl alcohol having an average of about 1 to about 5 ethylene oxide units, and mixtures thereof. More preferred structuring agents of the present invention are selected from stearyl alcohol, cetyl alcohol, behenyl alcohol, the polyethylene glycol ether of stearyl alcohol having an average of about 2 ethylene oxide units (steareth-2), the polyethylene glycol ether of stearyl alcohol having an average of about 21 ethylene oxide units (steareth-21),

the polyethylene glycol ether of cetyl alcohol having an average of about 2 ethylene oxide units, and mixtures thereof. Even more preferred structuring agents are selected from stearic acid, palmitic acid, stearyl alcohol, cetyl alcohol, behenyl alcohol, steareth-2, steareth-21, and mixtures thereof.

(2) Hydrophilic surfactant


The preferred oil-in-water emulsions contain from about 0.05% to about 10%, preferably from about 1% to about 6%, and more preferably from about 1% to about 3% of at least one hydrophilic surfactant which can disperse the hydrophobic materials in the water phase (percentages by weight of the topical carrier). The surfactant, at a minimum, must be hydrophilic enough to disperse in water.

Preferred hydrophilic surfactants are selected from nonionic surfactants. Among the nonionic surfactants that are useful herein are those that can be broadly defined as condensation products of long chain alcohols, e.g. C8-30 alcohols, with sugar or starch polymers, i.e., glycosides. These compounds can be represented by the formula $(S)_n-O-R$ wherein S is a sugar moiety such as glucose, fructose, mannose, and galactose; n is an integer of from about 1 to about 1000, and R is a C8-30 alkyl group. Examples of long chain alcohols from which the alkyl group can be derived include decyl alcohol, cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol, oleyl alcohol, and the like. Preferred examples of these surfactants include those wherein S is a glucose moiety, R is a C8-20 alkyl group, and n is an integer of from about 1 to about 9. Commercially available examples of these surfactants include decyl polyglucoside (available as APG 325 CS from Henkel) and lauryl polyglucoside (available as APG 600 CS and 625 CS from Henkel).

Other useful nonionic surfactants include the condensation products of alkylene oxides with fatty acids (i.e. alkylene oxide esters of fatty acids). These materials have the general formula $RCO(X)_nOH$ wherein R is a C10-30 alkyl group, X is $-OCH_2CH_2-$ (i.e. derived from ethylene glycol or oxide) or $-OCH_2CHCH_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 200. Other nonionic surfactants are the condensation products of alkylene oxides with 2 moles of fatty acids (i.e. alkylene oxide diesters of fatty acids). These materials have the general formula $RCO(X)_nOOCR$ wherein R is a C10-30 alkyl group, X is $-OCH_2CH_2-$ (i.e. derived from ethylene glycol or oxide) or $-OCH_2CHCH_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 100. Other nonionic surfactants are the condensation products of alkylene oxides with fatty alcohols (i.e.

alkylene oxide ethers of fatty alcohols). These materials have the general formula $R(X)_nOR'$ wherein R is a C10-30 alkyl group, X is $-OCH_2CH_2-$ (i.e. derived from ethylene glycol or oxide) or $-OCH_2CHCH_3-$ (i.e. derived from propylene glycol or oxide), and n is an integer from about 6 to about 100 and R' is H or a C10-30 alkyl group. Still other nonionic surfactants are the condensation products of alkylene oxides with both fatty acids and fatty alcohols [i.e. wherein the polyalkylene oxide portion is esterified on one end with a fatty acid and etherified (i.e. connected via an ether linkage) on the other end with a fatty alcohol]. These materials have the general formula $RCO(X)_nOR'$ wherein R and R' are C10-30 alkyl groups, X is $-OCH_2CH_2-$ (i.e. derived from ethylene glycol or oxide) or $-OCH_2CHCH_3-$ (derived from propylene glycol or oxide), and n is an integer from about 6 to about 100. Nonlimiting examples of these alkylene oxide derived nonionic surfactants include ceteth-6, ceteth-10, ceteth-12, cetareth-6, cetareth-10, cetareth-12, steareth-6, steareth-10, steareth-12, steareth-21, PEG-6 stearate, PEG-10 stearate, PEG-100 stearate, PEG-12 stearate, PEG-20 glyceryl stearate, PEG-80 glyceryl tallowate, PEG-10 glyceryl stearate, PEG-30 glyceryl cocoate, PEG-80 glyceryl cocoate, PEG-200 glyceryl tallowate, PEG-8 dilaurate, PEG-10 distearate, and mixtures thereof.

Still other useful nonionic surfactants include polyhydroxy fatty acid amide surfactants corresponding to the structural formula:



wherein: R^1 is H, C_1 - C_4 alkyl, 2-hydroxyethyl, 2-hydroxypropyl, preferably C_1 - C_4 alkyl, more preferably methyl or ethyl, most preferably methyl; R^2 is C_5 - C_{31} alkyl or alkenyl, preferably C_7 - C_{19} alkyl or alkenyl, more preferably C_9 - C_{17} alkyl or alkenyl, most preferably C_{11} - C_{15} alkyl or alkenyl; and Z is a polyhydroxyhydrocarbyl moiety having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxyated or propoxyated) thereof. Z preferably is a sugar moiety selected from the group consisting of glucose, fructose, maltose, lactose, galactose, mannose, xylose, and mixtures thereof. An especially preferred surfactant corresponding to the above structure is

coconut alkyl N-methyl glucoside amide (i.e., wherein the R^2CO- moiety is derived from coconut oil fatty acids). Processes for making compositions containing polyhydroxy fatty acid amides are disclosed, for example, in G.B. Patent Specification 809,060, published February 18, 1959, by Thomas Hedley & Co., Ltd.; U.S. Patent No. 2,965,576, to E. R. Wilson, issued December 20, 1960; U.S. Patent No. 2,703,798, to A. M. Schwartz, issued March 8, 1955; and U.S. Patent No. 1,985,424, to Piggott, issued December 25, 1934; which are incorporated herein by reference in their entirety.

Preferred among the nonionic surfactants are those selected from the group consisting of steareth-21, cetareth-20, cetareth-12, sucrose cocoate, steareth-100, PEG-100 stearate, and mixtures thereof.


Other nonionic surfactants suitable for use herein include sugar esters and polyesters, alkoxylated sugar esters and polyesters, C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxylated derivatives of C1-C30 fatty acid esters of C1-C30 fatty alcohols, alkoxylated ethers of C1-C30 fatty alcohols, polyglyceryl esters of C1-C30 fatty acids, C1-C30 esters of polyols, C1-C30 ethers of polyols, alkyl phosphates, polyoxyalkylene fatty ether phosphates, fatty acid amides, acyl lactylates, and mixtures thereof. Nonlimiting examples of these emulsifiers include: polyethylene glycol 20 sorbitan monolaurate (Polysorbate 20), polyethylene glycol 5 soya sterol, Steareth-20, Cetareth-20, PPG-2 methyl glucose ether distearate, Ceteth-10, Polysorbate 80, cetyl phosphate, potassium cetyl phosphate, diethanolamine cetyl phosphate, Polysorbate 60, glyceryl stearate, polyoxyethylene 20 sorbitan trioleate (Polysorbate 85), sorbitan monolaurate, polyoxyethylene 4 lauryl ether sodium stearate, polyglyceryl-4 isostearate, hexyl laurate, PPG-2 methyl glucose ether distearate, PEG-100 stearate, and mixtures thereof.

Another group of nonionic surfactants useful herein include the fatty acid ester blends based on a mixture of sorbitan or sorbitol fatty acid ester and sucrose fatty acid ester, the fatty acid in each instance being preferably C₈-C₂₄, more preferably C₁₀-C₂₀. The preferred fatty acid ester emulsifier is a blend of sorbitan or sorbitol C₁₆-C₂₀ fatty acid ester with sucrose C₁₀-C₁₆ fatty acid ester, especially sorbitan stearate and sucrose cocoate. This is commercially available from ICI under the trade name Arlatone 2121.

Other suitable surfactants useful herein include a wide variety of cationic, anionic, zwitterionic, and amphoteric surfactants such as are known in the art and discussed more fully below. See, e.g., McCutcheon's, Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation; U.S. Patent No. 5,011,681 to Ciotti et al., issued

April 30, 1991; U.S. Patent No. 4,421,769 to Dixon et al., issued December 20, 1983; and U.S. Patent No. 3,755,560 to Dickert et al., issued August 28, 1973; these four references are incorporated herein by reference in their entirety. The hydrophilic surfactants useful herein can contain a single surfactant, or any combination of suitable surfactants. The exact surfactant (or surfactants) chosen will depend upon the pH of the composition and the other components present.

The cationic surfactants useful herein include dialkyl quaternary ammonium compounds, examples of which are described in U.S. Patent 5,151,209; U.S. Patent 5,151,210; U.S. Patent 5,120,532; U.S. Patent 4,387,090; U.S. Patent 3,155,591; U.S. Patent 3,929,678; U.S. Patent 3,959,461; McCutcheon's, Detergents & Emulsifiers, (North American edition 1979) M.C. Publishing Co.; and Schwartz, et al., Surface Active Agents, Their Chemistry and Technology, New York: Interscience Publishers, 1949; which descriptions are incorporated herein by reference. The cationic surfactants useful herein also include cationic ammonium salts such as those having the formula:



wherein R_1 , is an alkyl group having from about 12 to about 30 carbon atoms, or an aromatic, aryl or alkaryl group having from about 12 to about 30 carbon atoms; R_2 , R_3 , and R_4 are independently selected from hydrogen, an alkyl group having from about 1 to about 22 carbon atoms, or aromatic, aryl or alkaryl groups having from about 12 to about 22 carbon atoms; and X is any compatible anion, preferably selected from chloride, bromide, iodide, acetate, phosphate, nitrate, sulfate, methyl sulfate, ethyl sulfate, tosylate, lactate, citrate, glycolate, and mixtures thereof. Additionally, the alkyl groups of R_1 , R_2 , R_3 , and R_4 can also contain ester and/or ether linkages, or hydroxy or amino group substituents (e.g., the alkyl groups can contain polyethylene glycol and polypropylene glycol moieties).

More preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 is selected from H or an alkyl group having from about 1 to about 22 carbon atoms; R_3 and R_4 are independently selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described previously.

Still more preferably, R_1 is an alkyl group having from about 12 to about 22 carbon atoms; R_2 , R_3 , and R_4 are selected from H or an alkyl group having from about 1 to about 3 carbon atoms; and X is as described previously.

Alternatively, other useful cationic emulsifiers include amino-amides, wherein in the above structure R_1 is alternatively $R_5\text{CONH}-(\text{CH}_2)_n$, wherein R_5 is an alkyl group having from about 12 to about 22 carbon atoms, and n is an integer from about 2 to about 6, more preferably from about 2 to about 4, and still more preferably from about 2 to about 3. Nonlimiting examples of these cationic emulsifiers include stearamidopropyl PG-dimonium chloride phosphate, behenamidopropyl PG dimonium chloride, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof. Especially preferred is behenamidopropyl PG dimonium chloride.

Nonlimiting examples of quaternary ammonium salt cationic surfactants include those selected from cetyl ammonium chloride, cetyl ammonium bromide, lauryl ammonium chloride, lauryl ammonium bromide, stearyl ammonium chloride, stearyl ammonium bromide, cetyl dimethyl ammonium chloride, cetyl dimethyl ammonium bromide, lauryl dimethyl ammonium chloride, lauryl dimethyl ammonium bromide, stearyl dimethyl ammonium chloride, stearyl dimethyl ammonium bromide, cetyl trimethyl ammonium chloride, cetyl trimethyl ammonium bromide, lauryl trimethyl ammonium chloride, lauryl trimethyl ammonium bromide, stearyl trimethyl ammonium chloride, stearyl trimethyl ammonium bromide, lauryl dimethyl ammonium chloride, stearyl dimethyl cetyl ditallow dimethyl ammonium chloride, dicetyl ammonium chloride, dicetyl ammonium bromide, dilauryl ammonium chloride, dilauryl ammonium bromide, distearyl ammonium chloride, distearyl ammonium bromide, dicetyl methyl ammonium chloride, dicetyl methyl ammonium bromide, dilauryl methyl ammonium chloride, dilauryl methyl ammonium bromide, distearyl methyl ammonium chloride, distearyl methyl ammonium bromide, and mixtures thereof. Additional quaternary ammonium salts include those wherein the C_{12} to C_{30} alkyl carbon chain is derived from a tallow fatty acid or from a coconut fatty acid. The term "tallow" refers to an alkyl group derived from tallow fatty acids (usually hydrogenated tallow fatty acids), which generally have mixtures of alkyl chains in the C_{16} to C_{18} range. The term "coconut" refers to an alkyl group derived from a coconut fatty acid, which generally have mixtures of alkyl chains in the C_{12} to C_{14} range. Examples of

quaternary ammonium salts derived from these tallow and coconut sources include ditallow dimethyl ammonium chloride, ditallow dimethyl ammonium methyl sulfate, di(hydrogenated tallow) dimethyl ammonium chloride, di(hydrogenated tallow) dimethyl ammonium acetate, ditallow dipropyl ammonium phosphate, ditallow dimethyl ammonium nitrate, di(coconutalkyl)dimethyl ammonium chloride, di(coconutalkyl)dimethyl ammonium bromide, tallow ammonium chloride, coconut ammonium chloride, stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldimonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof. An example of a quaternary ammonium compound having an alkyl group with an ester linkage is ditallowyl oxyethyl dimethyl ammonium chloride.

More preferred cationic surfactants are those selected from behenamidopropyl PG dimonium chloride, dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, stearamidopropyl PG-dimonium chloride phosphate, stearamidopropyl ethyldiammonium ethosulfate, stearamidopropyl dimethyl (myristyl acetate) ammonium chloride, stearamidopropyl dimethyl cetearyl ammonium tosylate, stearamidopropyl dimethyl ammonium chloride, stearamidopropyl dimethyl ammonium lactate, and mixtures thereof.


Still more preferred cationic surfactants are those selected from behenamidopropyl PG dimonium chloride, dilauryl dimethyl ammonium chloride, distearyl dimethyl ammonium chloride, dimyristyl dimethyl ammonium chloride, dipalmityl dimethyl ammonium chloride, and mixtures thereof.

A preferred combination of cationic surfactant and structuring agent is behenamidopropyl PG dimonium chloride and/or behenyl alcohol, wherein the ratio is preferably optimized to maintained to enhance physical and chemical stability, especially when such a combination contains ionic and/or highly polar solvents. This combination is especially useful for delivery of sunscreens agents such as zinc oxide and octyl methoxycinnamate.

A wide variety of anionic surfactants are also useful herein. See, e.g., U.S. Patent No. 3,929,678, to Laughlin et al., issued December 30, 1975, which is incorporated herein by reference in its entirety. Nonlimiting examples of anionic surfactants include the alkoyl isethionates, and the alkyl and alkyl ether sulfates. The alkoyl isethionates typically have the formula $\text{RCO-OCH}_2\text{CH}_2\text{SO}_3\text{M}$ wherein R is alkyl or alkenyl of from about 10 to about 30

carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Nonlimiting examples of these isethionates include those alkoyl isethionates selected from ammonium cocoyl isethionate, sodium cocoyl isethionate, sodium lauroyl isethionate, sodium stearyl isethionate, and mixtures thereof.

The alkyl and alkyl ether sulfates typically have the respective formulae ROSO_3M and $\text{RO}(\text{C}_2\text{H}_4\text{O})_x\text{SO}_3\text{M}$, wherein R is alkyl or alkenyl of from about 10 to about 30 carbon atoms, x is from about 1 to about 10, and M is a water-soluble cation such as ammonium, sodium, potassium and triethanolamine. Another suitable class of anionic surfactants are the water-soluble salts of the organic, sulfuric acid reaction products of the general formula:



wherein R_1 is chosen from the group including a straight or branched chain, saturated aliphatic hydrocarbon radical having from about 8 to about 24, preferably about 10 to about 16, carbon atoms; and M is a cation. Still other anionic synthetic surfactants include the class designated as succinamates, olefin sulfonates having about 12 to about 24 carbon atoms, and β -alkyloxy alkane sulfonates. Examples of these materials are sodium lauryl sulfate and ammonium lauryl sulfate.

Other anionic materials useful herein are soaps (i.e. alkali metal salts, e.g., sodium or potassium salts) of fatty acids, typically having from about 8 to about 24 carbon atoms, preferably from about 10 to about 20 carbon atoms. The fatty acids used in making the soaps can be obtained from natural sources such as, for instance, plant or animal-derived glycerides (e.g., palm oil, coconut oil, soybean oil, castor oil, tallow, lard, etc.) The fatty acids can also be synthetically prepared. Soaps are described in more detail in U.S. Patent No. 4,557,853.

Amphoteric and zwitterionic surfactants are also useful herein. Examples of amphoteric and zwitterionic surfactants which can be used in the compositions of the present invention are those which are broadly described as derivatives of aliphatic secondary and tertiary amines in which the aliphatic radical can be straight or branched chain and wherein one of the aliphatic substituents contains from about 8 to about 22 carbon atoms (preferably $\text{C}_8 - \text{C}_{18}$) and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Examples are alkyl imino acetates, and iminodialkanoates and aminoalkanoates of the formulas $\text{RN}(\text{CH}_2)_m\text{CO}_2\text{M}]_2$ and $\text{RNH}(\text{CH}_2)_m\text{CO}_2\text{M}$ wherein m is from 1 to 4, R is a $\text{C}_8\text{-C}_{22}$ alkyl or alkenyl, and M is H, alkali metal, alkaline earth metal ammonium, or

alkanolammonium. Also included are imidazolinium and ammonium derivatives. Specific examples of suitable amphoteric surfactants include sodium 3-dodecyl-aminopropionate, sodium 3-dodecylaminopropane sulfonate, N-alkyltaurines such as the one prepared by reacting dodecylamine with sodium isethionate according to the teaching of U.S. Patent 2,658,072 which is incorporated herein by reference in its entirety; N-higher alkyl aspartic acids such as those produced according to the teaching of U.S. Patent 2,438,091 which is incorporated herein by reference in its entirety; and the products sold under the trade name "Miranol" and described in U.S. Patent 2,528,378, which is incorporated herein by reference in its entirety. Other examples of useful amphoterics include phosphates, such as coamidopropyl PG-dimonium chloride phosphate (commercially available as Monaquat PTC, from Mona Corp.).

Other amphoteric or zwitterionic surfactants useful herein include betaines. Examples of betaines include the higher alkyl betaines, such as coco dimethyl carboxymethyl betaine, lauryl dimethyl carboxymethyl betaine, lauryl dimethyl alphacarboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, cetyl dimethyl betaine (available as Lonzaine 16SP from Lonza Corp.), lauryl bis-(2-hydroxyethyl) carboxymethyl betaine, stearyl bis-(2-hydroxypropyl) carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine, lauryl bis-(2-hydroxypropyl)alpha-carboxyethyl betaine, coco dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine, and amidobetaines and amidosulfobetaines (wherein the $\text{RCONH}(\text{CH}_2)_3$ radical is attached to the nitrogen atom of the betaine), oleyl betaine (available as amphoteric Velvetex OLB-50 from Henkel), and cocamidopropyl betaine (available as Velvetex BK-35 and BA-35 from Henkel).

Other useful amphoteric and zwitterionic surfactants include the sultaines and hydroxysultaines such as cocamidopropyl hydroxysultaine (available as Mirataine CBS from Rhone-Poulenc), and the alkanoyl sarcosinates corresponding to the formula $\text{RCON}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CO}_2\text{M}$ wherein R is alkyl or alkenyl of about 10 to about 20 carbon atoms, and M is a water-soluble cation such as ammonium, sodium, potassium and trialkanolamine (e.g., triethanolamine), a preferred example of which is sodium lauroyl sarcosinate.

(3) Water

The preferred oil-in-water emulsion contains from about 25% to about 98%, preferably from about 65% to about 95%, more preferably from about 70% to about 90% water by weight of the topical carrier.

The hydrophobic phase is dispersed in the continuous aqueous phase. The hydrophobic phase may contain water insoluble or partially soluble materials such as are known in the art, including but not limited to the silicones described herein in reference to silicone-in-water emulsions, and other oils and lipids such as described above in reference to emulsions.

The topical compositions of the subject invention, including but not limited to lotions and creams, may contain a dermatologically acceptable emollient. Such compositions preferably contain from about 1% to about 50% of the emollient. As used herein, "emollient" refers to a material useful for the prevention or relief of dryness, as well as for the protection of the skin. A wide variety of suitable emollients are known and may be used herein. Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 32-43 (1972), incorporated herein by reference, contains numerous examples of materials suitable as an emollient. A preferred emollient is glycerin. Glycerin is preferably used in an amount of from or about 0.001 to or about 30%, more preferably from or about 0.01 to or about 20%, still more preferably from or about 0.1 to or about 10%, e.g., 5%.

Lotions and creams according to the present invention generally contain a solution carrier system and one or more emollients. Lotions and creams typically contain from about 1% to about 50%, preferably from about 1% to about 20%, of emollient; from about 50% to about 90%, preferably from about 60% to about 80%, water; and farnesol, bisabolol and the additional skin care active in the above described amounts. Creams are generally thicker than lotions due to higher levels of emollients or higher levels of thickeners.

Ointments of the present invention may contain a simple carrier base of animal or vegetable oils or semi-solid hydrocarbons (oleaginous); absorption ointment bases which absorb water to form emulsions; or water soluble carriers, e.g., a water soluble solution carrier. Ointments may further contain a thickening agent, such as described in Sagarin, Cosmetics, Science and Technology, 2nd Edition, Vol. 1, pp. 72-73 (1972), incorporated herein by reference, and/or an emollient. For example, an ointment may contain from about 2% to about 10% of an emollient; from about 0.1% to about 2% of a thickening agent; and farnesol, bisabolol and the additional skin care active in the above described amounts.

Compositions of this invention useful for cleansing ("cleansers") are formulated with a suitable carrier, e.g., as described above, and preferably contain, in addition to farnesol, bisabolol and the additional skin care active in the above described amounts, from about 1% to about 90%, more preferably from about 5% to about 10%, of a dermatologically acceptable surfactant. The surfactant is suitably selected from anionic, nonionic, zwitterionic, amphoteric

and ampholytic surfactants, as well as mixtures of these surfactants. Such surfactants are well known to those skilled in the detergency art. Nonlimiting examples of possible surfactants include isoceteth-20, sodium methyl cocoyl taurate, sodium methyl oleoyl taurate, and sodium lauryl sulfate. See U.S. Patent No. 4,800,197, to Kowcz et al., issued January 24, 1989, which is incorporated herein by reference in its entirety, for exemplary surfactants useful herein. Examples of a broad variety of additional surfactants useful herein are described in McCutcheon's Detergents and Emulsifiers, North American Edition (1986), published by Allured Publishing Corporation. The cleansing compositions can optionally contain, at their art-established levels, other materials which are conventionally used in cleansing compositions.

The physical form of the cleansing compositions is not critical. The compositions can be, for example, formulated as toilet bars, liquids, shampoos, bath gels, hair conditioners, hair tonics, pastes, or mousses. Rinse-off cleansing compositions, such as shampoos, require a delivery system adequate to deposit sufficient levels of actives on the skin and scalp. A preferred delivery system involves the use of insoluble complexes. For a more complete disclosure of such delivery systems, see U.S. Patent 4,835,148, Barford et al., issued May 30, 1989.

As used herein, the term "foundation" refers to a liquid, semi-liquid, semi-solid, or solid skin cosmetic which includes, but is not limited to lotions, creams, gels, pastes, cakes, and the like. Typically the foundation is used over a large area of the skin, such as over the face, to provide a particular look. Foundations are typically used to provide an adherent base for color cosmetics such as rouge, blusher, powder and the like, and tend to hide skin imperfections and impart a smooth, even appearance to the skin. Foundations of the present invention include a dermatologically acceptable carrier and may include conventional ingredients such as oils, colorants, pigments, emollients, fragrances, waxes, stabilizers, and the like. Exemplary carriers and such other ingredients which are suitable for use herein are described, for example, in PCT Application, WO 96/33689, to Canter, et al., published on October 31, 1996 and U.K. Patent, GB 2274585, issued on August 3, 1994.

Composition Preparation

The compositions useful for the methods of the present invention are generally prepared by conventional methods such as are known in the art of making topical compositions. Such methods typically involve mixing of the ingredients in one or more steps to a relatively uniform state, with or without heating, cooling, application of vacuum, and the like.

Methods for Regulating Skin Condition

The compositions of the present invention are useful for regulating mammalian skin condition. Such regulation of keratinous tissue conditions can include prophylactic and therapeutic regulation. For example, such regulating methods are directed to thickening keratinous tissue (i.e., building the epidermis and/or dermis layers of the skin and where applicable the keratinous layers of the nail and hair shaft) and preventing and/or retarding atrophy of mammalian skin, preventing and/or retarding the appearance of spider vessels and/or red blotchiness on mammalian skin, preventing and/or retarding the appearance of dark circles under the eye of a mammal, preventing and/or retarding sallowness of mammalian skin, preventing and/or retarding sagging of mammalian skin, softening and/or smoothing lips, hair and nails of a mammal, preventing and/or relieving itch of mammalian skin, regulating skin texture (e.g. wrinkles and fine lines), and improving skin color (e.g. redness, freckles).

Regulating keratinous tissue condition involves topically applying to the keratinous tissue a safe and effective amount of a composition of the present invention. The amount of the composition which is applied, the frequency of application and the period of use will vary widely depending upon the level of farnesol, bisabolol and the additional skin care active and/or other components of a given composition and the level of regulation desired, e.g., in light of the level of keratinous tissue damage present or expected to occur.

In a preferred embodiment, the composition is chronically applied to the skin. By "chronic topical application" is meant continued topical application of the composition over an extended period during the subject's lifetime, preferably for a period of at least about one week, more preferably for a period of at least about one month, even more preferably for at least about three months, even more preferably for at least about six months, and more preferably still for at least about one year. While benefits are obtainable after various maximum periods of use (e.g., five, ten or twenty years), it is preferred that chronic application continue throughout the subject's lifetime. Typically applications would be on the order of about once per day over such extended periods, however application rates can vary from about once per week up to about three times per day or more.

A wide range of quantities of the compositions of the present invention can be employed to provide a skin appearance and/or feel benefit. Quantities of the present compositions which are typically applied per application are, in mg composition/cm² skin, from about 0.1 mg/cm² to about 10 mg/cm². A particularly useful application amount is about 1 mg/cm² to about 2 mg/cm².

Regulating keratinous tissue condition is preferably practiced by applying a composition in the form of a skin lotion, cream, gel, foam, ointment, paste, emulsion, spray, conditioner, tonic, cosmetic, lipstick, foundation, nail polish, after-shave, or the like which is preferably intended to be left on the skin or other keratin structure for some esthetic, prophylactic, therapeutic or other benefit (i.e., a "leave-on" composition). After applying the composition to the skin, it is preferably left on the skin for a period of at least about 15 minutes, more preferably at least about 30 minutes, even more preferably at least about 1 hour, still more preferably for at least several hours, e.g., up to about 12 hours. Any part of the external portion of the face, hair, and/or nails can be treated, e.g., face, lips, under-eye area, eyelids, scalp, neck, torso, arms, hands, legs, feet, fingernails, toenails, scalp hair, eyelashes, eyebrows, etc. The composition can be applied with the fingers or with an implement or device (e.g., pad, cotton ball, applicator pen, spray applicator, and the like).

Another approach to ensure a continuous exposure of the skin to at least a minimum level of the farnesol, bisabolol and the additional skin care active is to apply the compound by use of a patch applied, e.g., to the face. Such an approach is particularly useful for problem skin areas needing more intensive treatment (e.g., facial crows feet area, frown lines, under eye area, and the like). The patch can be occlusive, semi-occlusive or non-occlusive and can be adhesive or non-adhesive. The farnesol, bisabolol and the additional skin care active composition can be contained within the patch or be applied to the skin prior to application of the patch. The patch can also include additional actives such as chemical initiators for exothermic reactions such as those described in U.S. Patents numbered 5,821,250, 5,981,547, and 5,972,957 to Wu, et al. The patch is preferably left on the skin for a period of at least about 5 minutes, more preferably at least about 15 minutes, more preferably still at least about 30 minutes, even more preferably at least about 1 hour, still more preferably at night as a form of night therapy.

Examples

The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

Examples 1 - 6

A skin cream is prepared by conventional methods from the following components.

	Ingredient	Ex 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6
PHASE A:	Water U.S.P.	qs to 100	qs to 100	qs to 100	qs to 100	qs to 100	qs to 100

	Disodium EDTA	0.1	0.1	0.1	0.1	0.1	0.1
	Methyl Paraben	0.25	0.25	0.25	0.25	0.25	0.25
	Niacinamide					2.0	3.5
	Dexpantenol					1.0	1.0
	Allantoin	0.20	0.20	0.20	0.20	0.20	0.20
	Glycerin	7.0	7.0	7.0	7.0	7.0	7.0
PHASE B:	Cetyl Alcohol	0.3	0.3	0.3	0.3	0.3	0.3
	Stearyl Alcohol	0.5	0.5	0.5	0.5	0.5	0.5
	Behenyl Alcohol	0.4	0.4	0.4	0.4	0.4	0.4
	Vitamin E Acetate					0.5	0.5
	Mryj 59	0.10	0.10	0.10	0.10	0.10	0.10
	Propyl Paraben	0.10	0.10	0.10	0.10	0.10	0.10
	Farnesol	5.0	5.0	5.0	5.0	5.0	5.0
	(-)-alpha-Bisabolol	1.0	1.0	1.0	1.0	1.0	1.0
	Phytantriol	5.0			1.0		
	Permethyl 101 A	3.3	3.0	3.0	3.0	3.0	3.0
PHASE C	Sepigel 305	2.0	2.0	2.0	2.0	2.0	2.0
PHASE D	Titanium Dioxide GLW75CA*		0.5				
PHASE E	Benzyl Alcohol	0.50	0.50	0.50	0.50	0.50	0.50
	Dimethicone/Dimethiconol	0.50	0.50	0.50	0.50	0.50	0.50
PHASE F	Palmitoyl-Lys-Thr-Thr-Lys-Ser ¹			3.0	3.0		3.0
PHASE G:	Fragrance	0.20	0.20	0.20	0.20	0.20	0.20

¹ Peptide can be obtained from Sederma as Matrixyl (100 ppm of palmitoyl-Lys-Thr-Thr-Lys-Ser)

* Available from KOBO

Blend the A phase components with a suitable mixer (e.g., Tekmar model RW20DZM), heating while stirring to a temperature of 70-80°C. Separately, blend the B phase components with a suitable mixer and heat to 70 - 75 C and maintain while mixing. Phase B is added to Phase A while mixing well to emulsify. When emulsion is at approx. 60 C, Phase C is added while continuing to mix emulsion. The emulsion is allowed to cool to approx. 40 C while stirring. At approx. 50 C, Phase D and E are added to the emulsion and mixing continued. At approx. 40 C, Phases F and G are added to the emulsion. The emulsion is then milled using a suitable mill (Tekmar T-25) for approx. 5 minutes resulting in an uniform product.

Examples 7 - 10

A skin lotion is prepared by conventional methods from the following components.

	Ingredient	Ex 7	Ex. 8	Ex. 9	Ex. 10
PHASE A	Water U.S.P.	qs to 100	qs to 100	qs to 100	qs to 100
	Disodium EDTA	0.10	0.10	0.10	0.10
	Allantoin	0.20	0.20	0.20	0.20
	Glycerin	5.0	5.0	5.0	5.0
	Niacinamide				3.5
PHASE B	Cetyl Alcohol	1.2	1.2	1.2	1.2
	Stearyl Alcohol	1.2	1.2	1.2	1.2
	Brij 721	0.5	0.5	0.5	0.5
	Brij 72	0.1	0.1	0.1	0.1
	Salicylic Acid	1.5	1.5	1.5	1.5
	PPG-15 Stearyl Ether	4.0	4.0	4.0	4.0
	Farnesol	3.0	3.0	3.0	3.0
	(-)-alpha-Bisabolol	1.0	1.0	0.5	1.0
	Phytantriol				
	Vitamin E Acetate			0.5	
	Titanium Dioxide SAT-T-CR-50			0.5	0.50
PHASE C	Sepigel 305	1.0	1.0	1.0	1.0
PHASE D	Dimethicone/Dimethiconol	0.5	0.5	0.5	0.5
PHASE E	Palmitoyl-Lys-Thr-Thr-Lys-Ser ¹		3.0	3.0	
PHASE F	Fragrance	0.20	0.20	0.20	0.20

¹ Peptide can be obtained from Sederma as Matrixyl (100 ppm of (palmitoyl-Lys-Thr-Thr-Lys-Ser)

* Available from U.S. Cosmetics

Blend the A phase components with a suitable mixer (e.g., Tekmar model RW20DZM), heating while stirring to a temperature of 70-80°C. Separately, blend the B phase components with a suitable mixer and heat to 70 - 75°C and maintain while mixing. Phase B is added to Phase A while mixing well to emulsify. When emulsion is at approx. 60°C, Phase C is added while continuing to mix emulsion. The emulsion is allowed to cool to approx. 40°C while stirring. At approx. 50°C, Phase D is added to the emulsion and mixing continued. At approx. 40°C, Phases E and F are added to the emulsion. The emulsion is then milled using a suitable mill (Tekmar T-25) for approx. 5 minutes resulting in an uniform product.

Examples 12 - 17

A skin cream is prepared by conventional methods from the following components.

	Ingredient	Ex 12	Ex. 13	Ex. 14	Ex. 15	Ex. 16	Ex. 17
PHASE A:	Water U.S.P.	qs to 100	qs to 100	qs to 100	qs to 100	qs to 100	qs to 100
	Disodium EDTA	0.15	0.15	0.15	0.15	0.15	0.15
	Methyl Paraben	0.25	0.25	0.25	0.25	0.25	0.25
	Allantoin	0.20	0.20	0.20	0.20	0.20	0.20
	Dexpanthenol						1.0
	Glycerin	3.0	3.0	3.0	3.0	3.0	5.0
PHASE B:	Cetyl Alcohol	0.30	0.30	0.30	0.30	0.30	0.30
	Stearyl Alcohol	0.50	0.50	0.50	0.50	0.50	0.50
	Behenyl Alcohol	0.40	0.40	0.40	0.40	0.40	0.40
	Myrj 59	0.10	0.10	0.10	0.10	0.10	0.10
	Propyl Paraben	0.10	0.10	0.10	0.10	0.10	0.10
	Farnesol	5.0	5.0	3.0	3.0	5.0	5.0
	(-)-alpha-Bisabolol	1.0	0.5	1.0	0.5	1.0	1.0
	Permethyl 101 A	3.25	3.25	3.25	3.25	3.25	3.25
	Octyl methoxycinnamate	1.5	2.0	7.0	6.0		
	4,4'-t-butyl methoxydibenzoylmethane	2.0	2.0			2.0	
	Octocrylene		5.0			1.5	
	Zinc Oxide				3.0		
	Terephthalylidene Dicamphor Sulfonic Acid			3.0			
	Octyl Salicylate					4.0	
PHASE C	Sepigel 305	2.0	2.0	2.0	2.0	2.0	2.0
PHASE D	Titanium Dioxide GLW75CA*	0.5		0.5			0.50
PHASE E	Benzyl Alcohol	0.50	0.50	0.50	0.50	0.50	0.50
	Dimethicone/Dimethiconol	0.75	0.75	0.75	0.75	0.75	0.75
PHASE F	Palmitoyl-Lys-Thr-Thr-Lys-Ser ¹			3.0			
PHASE G	Water						2.0
	Finsolv TN						1.5
	Polysorbate -20						0.20
	Retinyl propionate						0.20
PHASE H	Fragrance	0.10	0.10	0.10	0.10	0.20	0.20

¹ Peptide can be obtained from Sederma as Matrixyl (100 ppm of palmitoyl-Lys-Thr-Thr-Lys-Ser)

* Available from KOBO

Blend the A phase components with a suitable mixer (e.g., Tekmar model RW20DZM), heating while stirring to a temperature of 70-80°C. Separately, blend the B phase components

with a suitable mixer and heat to 70 - 75°C and maintain while mixing. Phase B is added to Phase A while mixing well to emulsify. When emulsion is at approx. 60°C, Phase C is added while continuing to mix emulsion. The emulsion is allowed to cool to approx. 40°C while stirring. At approx. 50°C, Phase D and E are added to the emulsion and mixing continued. At approx. 40°C, Phases F, G and H are added to the emulsion. The emulsion is then milled using a suitable mill (Tekmar T-25) for approx. 5 minutes resulting in a uniform product.

WHAT IS CLAIMED IS:

1. A skin care composition characterized in that the composition comprises
 - a) a safe and effective amount, preferably from 0.1% to 10%, by weight of the composition, of farnesol;
 - b) a safe and effective amount, preferably from 0.1% to 10%, by weight of the composition, of bisabolol;
 - c) a safe and effective amount of at least one additional skin care active; and
 - d) a dermatologically acceptable carrier.
2. A skin care composition according to Claim 1, wherein the additional skin care active is selected from the group consisting of desquamatory actives, anti-acne actives, vitamin B₃ compounds, retinoids, peptides, hydroxy acids, anti-oxidants, radical scavengers, chelators, anti-inflammatory agents, topical anesthetics, tanning actives, skin lightening agents, anti-cellulite agents, flavonoids, antimicrobial actives, skin soothing agents, skin healing agents, antifungal actives, sunscreen actives, conditioning agents, structuring agents, thickening agents, and mixtures thereof.
3. A skin care composition according to Claim 1, wherein the additional skin care active is phytantriol.
4. A skin care composition according to Claim 1 or 3, wherein the additional skin care active is selected from the group consisting of vitamin B₃ compounds, and mixtures thereof, preferably the additional skin care active is niacinamide.
5. A skin care composition according to Claim 1, 3, or 4, wherein the additional skin care active is salicylic acid.
6. A skin care composition according to Claim 1, 3, 4, or 5, wherein the additional skin care active is a peptide active selected from the group consisting of lys-thr-thr-lys-ser, gly-his-lys, beta-ala-his, arg-ser-arg-lys, arg-lys-arg, his-gly-gly, their derivatives, and mixtures thereof, preferably the peptide active is palmitoyl-lys-thr-thr-lys-ser.
7. A skin care composition according to Claim 1, wherein the additional skin care active is a mixture of niacinamide, tocopherol acetate, and dexpanthenol.
8. A skin care composition according to Claim 1 or 2, wherein the additional skin care active is an inorganic sunscreen comprising a metallic oxide selected from the group consisting of titanium dioxide having an average primary particle size of from about 15 nm to about 100 nm, zinc oxide having an average primary particle size of from about 15 nm to about 150 nm, zirconium oxide having an average primary particle size of from about 15 nm to about 150 nm,

iron oxide having an average primary particle size of from about 15 nm to about 500nm, and mixtures thereof.

9. A skin care composition according to Claim 1 or 2, wherein the additional skin care active is an organic sunscreen selected from the group consisting of octylmethoxycinnamate, octyl salicylate, terephthalydene dicamphor sulfonic acid, avobenzone, octocrylene, and mixtures thereof.

10. Use of a skin care composition according to any one of the preceding claims to make a medicament for regulating the condition of mammalian skin by topical application to the skin of mammal in need of treatment.

SEQUENCE LISTING

<110> Robinson, Larry R.
Bissett, Donald L.
Deckner, George E.
Ha, Robert B.K.

<120> Skin Care Compositions Containing Combination of Skin
Care Actives

<130> skin care composition 3

<140> 8023

<141> 2000-04-07

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
palmitoyl-lys-thr-thr-lys-ser

<400> 1

Lys Thr Thr Lys Ser
1 5

<210> 2

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
arg-ser-arg-lys

<400> 2

Arg Ser Arg Lys
1